



United  
Healthcare®  
Community Plan

UNITEDHEALTHCARE® COMMUNITY PLAN:  
RADIOLOGY IMAGING COVERAGE DETERMINATION GUIDELINE

## Adult Musculoskeletal Imaging Guidelines (For Ohio Only)

**V2.0.2024**

Guideline Number: CSRAD007OH.C

*Effective Date: November 15, 2024*

### Application (for Ohio Only)

*This Medical Policy only applies to the state of Ohio. Any requests for services that are stated as unproven or services for which there is a coverage or quantity limit will be evaluated for medical necessity using Ohio Administrative Code 5160-1-01.*

# Table of Contents

## Guideline

---

### **Related Community Plan Policies**

#### **Application (For Ohio Only)**

#### **Guideline Development (Preface-1)**

#### **Benefits, Coverage Policies, and Eligibility Issues (Preface-2)**

#### **Clinical Information (Preface-3)**

#### **Coding Issues (Preface-4)**

#### **Whole-Body Imaging (Preface-5)**

#### **References (Preface-6)**

#### **Copyright Information (Preface-7)**

#### **Trademarks (Preface-8)**

#### **General Guidelines (MS-1)**

#### **Imaging Techniques (MS-2)**

#### **3D Rendering (MS-3)**

#### **Avascular Necrosis (AVN)/Osteonecrosis (MS-4)**

#### **Fractures (MS-5)**

#### **Foreign Body (MS-6)**

#### **Ganglion Cysts (MS-7)**

#### **Gout/Calcium Pyrophosphate Deposition Disease (CPPD)/ Pseudogout/**

#### **Chondrocalcinosis (MS-8)**

#### **Infection/Osteomyelitis (MS-9)**

#### **Soft Tissue Mass or Lesion of Bone (MS-10)**

#### **Muscle/Tendon Unit Injuries/Diseases (MS-11)**

#### **Osteoarthritis (MS-12)**

#### **Chondral/Osteochondral Lesions (MS-13)**

#### **Osteoporosis (MS-14)**

#### **Rheumatoid Arthritis (RA) and Inflammatory Arthritis (MS-15)**

#### **Post-Operative Joint Replacement Surgery (MS-16)**

#### **Limb Length Discrepancy (MS-17)**

#### **Anatomical Area Tables – General Information (MS-18)**

#### **Shoulder (MS-19)**

#### **Elbow (MS-20)**

#### **Wrist (MS-21)**

#### **Hand (MS-22)**

#### **Pelvis (MS-23)**

#### **Hip (MS-24)**

#### **Knee (MS-25)**

**Ankle (MS-26)**  
**Foot (MS-27)**  
**Nuclear Medicine (MS-28)**  
**Policy History and Instructions for Use**

# Related Community Plan Policies

---

## Guideline

---

Related Community Plan Policies

# Related Community Plan Policies

## Related Community Plan Policies v2.0.2024

### General Policies

- Spine Imaging Guidelines
- Peripheral Vascular Disease (PVD) Imaging Guidelines
- Peripheral Nerve Disorders (PND) Imaging Guidelines

### Pediatric Policies

- Pediatric Musculoskeletal Imaging Guidelines

# Application (For Ohio Only)

---

## Guideline

---

Application (For Ohio Only)

# Application (For Ohio Only)

---

## Application for Ohio OH UHC v2.0.2024

- This Medical Policy only applies to the state of Ohio. Any requests for services that are stated as unproven or services for which there is a coverage or quantity limit will be evaluated for medical necessity using Ohio Administrative Code 5160-1-01.

# Guideline Development (Preface-1)

---

## Guideline

---

Guideline Development (Preface-1.1)



# Guideline Development (Preface-1.1)

PRF.GG.0001.1.UOH

v2.0.2024

- The UnitedHealthcare's evidence-based, proprietary clinical guidelines evaluate a range of advanced imaging and procedures, including NM, US, CT, MRI, PET, Radiation Oncology, Sleep Studies, as well as Cardiac, musculoskeletal and Spine interventions.
- UnitedHealthcare reserves the right to change and update the guidelines. The guidelines undergo a formal review annually. UnitedHealthcare's guidelines are based on current evidence supported by major national and international association and society guidelines and criteria, peer-reviewed literature, major treatises as well as, input from health plans, and practicing academic and community-based physicians.
- These guidelines are not intended to supersede or replace sound medical judgment, but instead, should facilitate the identification of the most appropriate imaging or other designated procedure given the individual's clinical condition. These guidelines are written to cover medical conditions as experienced by the majority of individuals. However, these guidelines may not be applicable in certain clinical circumstances, and physician judgment can override the guidelines.
- These guidelines provide evidence-based, clinical benefits with a focus on health care quality and patient safety.
- Clinical decisions, including treatment decisions, are the responsibility of the individual and his/her provider. Clinicians are expected to use independent medical judgment, which takes into account the clinical circumstances to determine individual management decisions.
- UnitedHealthcare supports the Choosing Wisely initiative (<https://www.choosingwisely.org/>) by the American Board of Internal Medicine (ABIM) Foundation and many national physician organizations, to reduce the overuse of diagnostic tests that are low value, no value, or whose risks are greater than the benefits.

# Benefits, Coverage Policies, and Eligibility Issues (Preface-2)

---

## Guideline

---

Benefits, Coverage Policies, and Eligibility Issues (Preface-2.1)  
References (Preface-2)

# Benefits, Coverage Policies, and Eligibility Issues (Preface-2.1)

PRF.BC.0002.1.UOH

v2.0.2024

## Investigational and Experimental Studies

- Certain studies, treatments, procedures, or devices may be considered experimental, investigational, or unproven for any condition, illness, disease, injury being treated if one of the following is present:
  - if there is a paucity of supporting evidence;
  - if the evidence has not matured to exhibit improved health parameters;
  - if clinical utility has not been demonstrated in any condition; OR
  - if the study, treatment, procedure, or device lacks a collective opinion of support
- Supporting evidence includes standards that are based on credible scientific evidence published in peer-reviewed medical literature (such as well conducted randomized clinical trials or cohort studies with a sample size of sufficient statistical power) generally recognized by the relevant medical community. Collective opinion of support includes physician specialty society recommendations and the views of physicians practicing in relevant clinical areas when physician specialty society recommendations are not available.

## Clinical and Research Trials

- Similar to investigational and experimental studies, clinical trial imaging requests will be considered to determine whether they meet UnitedHealthcare's evidence-based guidelines.
- Imaging studies which are inconsistent with established clinical standards, or are requested for data collection and not used in direct clinical management are not supported.

## Legislative Mandate

- State and federal legislations may need to be considered in the review of advanced imaging requests.

---

## References (Preface-2)

---

**v2.0.2024**

1. Coverage of Clinical Trials under the Patient Protection and Affordable Care Act; 42 U.S.C.A. § 300gg-8.

# Clinical Information (Preface-3)

---

## Guideline

---

Clinical Information (Preface-3.1)

References (Preface-3)

# Clinical Information (Preface-3.1)

PRF.CL.0003.1.UOH

v2.0.2024

## Clinical Documentation and Age Considerations

- UnitedHealthcare's guidelines use an evidence-based approach to determine the most appropriate procedure for each individual, at the most appropriate time in the diagnostic and treatment cycle. UnitedHealthcare's guidelines are framed by:
  - Clinical presentation of the individual, rather than the studies requested
  - Adequate clinical information that must be submitted to UnitedHealthcare in order to establish medical necessity for advanced imaging or other designated procedures includes, but is not limited to, the following:
    - Pertinent clinical evaluation should include a recent detailed history, physical examination<sup>20</sup> since the onset or change in symptoms, and/or laboratory and prior imaging studies.
      - Condition-specific guideline sections may describe additional clinical information which is required for a pertinent clinical evaluation.
      - The Spine and Musculoskeletal guidelines require x-ray studies from when the current episode of symptoms has started or changed; x-ray imaging does not have to be within the past 60 days.
      - Advanced imaging or other designated procedures should not be ordered prior to clinical evaluation of an individual by the physician treating the individual. This may include referral to a consultant specialist who will make further treatment decisions.
      - Other meaningful technological contact (telehealth visit, telephone or video call, electronic mail or messaging) since the onset or change in symptoms by an established individual can serve as a pertinent clinical evaluation.
        - Some conditions may require a face-to-face evaluation as discussed in the applicable condition-specific guideline sections.
    - A recent clinical evaluation may be unnecessary if the individual is undergoing a guideline-supported, scheduled follow-up imaging or other designated procedural evaluation. Exceptions due to routine surveillance indications are addressed in the applicable condition-specific guideline sections.
  - UnitedHealthcare's evidence-based approach to determine the most appropriate procedure for each individual requires submission of medical records pertinent to the requested imaging or other designated procedures.
- Many conditions affecting the pediatric population are different diagnoses than those occurring in the adult population. For those diseases which occur in both pediatric and adult populations, minor differences may exist in management due to individual

age, comorbidities, and differences in disease natural history between children and adults.

- Individuals who are 18 years old or younger<sup>19</sup> should be imaged according to the Pediatric Imaging Guidelines if discussed in the condition-specific guideline sections. Any conditions not specifically discussed in the Pediatric Imaging Guidelines should be imaged according to the General Imaging Guidelines. Individuals who are >18 years old should be imaged according to the General Imaging Guidelines, except where directed otherwise by a specific guideline section.
- The terms “male” and “female” used in these guidelines refer to anatomic-specific diseases and disease predispositions associated with the individual's sex assigned at birth rather than their gender identity. It should be noted that gender identity and anatomic-specific diseases as well as disease predispositions are not always linked. As such, these guidelines should be applied to the individual's corresponding known or suspected anatomic-specific disease or disease predisposition. At UnitedHealthcare, we believe that it is important to understand how all individuals, including those who are gender-diverse, choose to identify themselves. To ensure that gender-diverse individuals are treated with respect and that decisions impacting their healthcare are made correctly and with sensitivity, UnitedHealthcare recognizes all individuals with the following gender marker options: Male, Female, Transgender Male, Transgender Female, “X”, and “Not Specified.”

### **General Imaging Information**

- “Standard” or “conventional” imaging is most often performed in the initial and subsequent evaluations of malignancy. Standard or conventional imaging includes plain film, CT, MRI, or US.
  - Often, further advanced imaging is needed when initial imaging, such as ultrasound, CT, or MRI does not answer the clinical question. Uncertain, indeterminate, inconclusive, or equivocal may describe these situations.
- Appropriate use of contrast is a very important component of evidence-based advanced imaging use.
  - The appropriate levels of contrast for an examination (i.e., without contrast, with contrast, without and with contrast) is determined by the evidence-based guidance reflected in the condition-specific guideline sections.
  - If, during the performance of a non-contrast imaging study, there is the unexpected need to use contrast in order to evaluate a possible abnormality, then that is appropriate.<sup>1</sup>

### **Ultrasound**

- Diagnostic ultrasound uses high-frequency sound waves to evaluate soft tissue structures and vascular structures utilizing grey scale and Doppler techniques.
- Ultrasound allows for dynamic real-time imaging at the bedside.

- Ultrasound is limited in areas where there is dense bone or other calcification.
- Ultrasound also has a relatively limited imaging window so may be of limited value in evaluating very large abnormalities.
- In general, ultrasound is highly operator-dependent, and proper training and experience are required to perform consistent, high-quality evaluations.
- Indications for ultrasound may include, but are not limited to, the following:
  - Obstetric and gynecologic imaging
  - Soft tissue and visceral imaging of the chest, abdomen, pelvis, and extremities
  - Brain and spine imaging when not obscured by dense bony structures
  - Vascular imaging when not obscured by dense bony structures
  - Procedural guidance when not obscured by dense bony structures
  - Initial evaluation of ill-defined soft tissue masses or fullness and differentiating adenopathy from mass or cyst. Prior to advanced imaging, ultrasound can be very beneficial in selecting the proper modality, body area, image sequences, and contrast level that will provide the most definitive information for the individual.
- More specific guidance for ultrasound usage, including exceptions to this general guidance, can be found throughout the condition-specific guidelines.

### **Computed Tomography (CT)**

- The AMA CPT<sup>®</sup> manual does not describe nor assign any minimum or maximum number of sequences for any CT study. CT imaging protocols are often influenced by the individual's clinical situation and additional sequences are not uncommon. There are numerous CT protocols that may be performed to evaluate specific clinical questions, and this technology is constantly undergoing development.
- CT utilizes ionizing radiation to create cross-sectional and volumetric images of the body.
  - Advantages over ultrasound include a much larger field of view and faster completion time in general. Disadvantages compared to ultrasound include lack of portability and exposure to ionizing radiation.
  - Advantages over MRI include faster imaging and a more spacious scanner area limiting claustrophobia. Disadvantages compared to MRI include decreased soft tissue definition, especially with non-contrast imaging, and exposure to ionizing radiation.
- CT can be performed without, with, or without and with intravenous (IV) contrast depending on the clinical indication and body area.
  - In general, non-contrast imaging is appropriate for evaluating structures with significant tissue density differences such as lung parenchyma and bony structures, or when there is a contraindication to contrast.
  - In general, CT with contrast is the most common level of contrast and can be used when there is need for improved vascular or soft tissue resolution, including better



- characterization of known or suspected malignancy, as well as infectious and inflammatory conditions.
- CT without and with contrast has a limited role as the risks of doubling the ionizing radiation exposure rarely outweigh the benefits of multiphasic imaging, though there are some exceptions which include, but are not limited to, the following:
    - Characterization of a mass
    - Characterization of arterial and venous anatomy
    - CT with contrast may be used to better characterize findings on a very recent (within two weeks) inconclusive non-contrast CT where the guidelines would support CT without and with contrast.
  - More specific guidance for CT contrast usage, including exceptions to this general guidance, can be found throughout the condition-specific guidelines.
  - Shellfish allergy:
    - It is commonly assumed that an allergy to shellfish indicates iodine allergy, and that this implies an allergy to iodinated contrast media used with CT. However, this is NOT true. Shellfish allergy is due to tropomyosins. Iodine plays no role in these allergic reactions. Allergies to shellfish do not increase the risk of reaction to iodinated contrast media any more than that of other allergens.<sup>1</sup>
  - Enteric contrast (oral or rectal) is sometimes used in abdominal imaging. There is no specific CPT<sup>®</sup> code which refers to enteric contrast.
  - The appropriate contrast level and anatomic region in CT imaging is specific to the clinical indication, as listed in the condition-specific guideline sections.
  - CT should not be used to replace MRI in an attempt to avoid sedation unless it is listed as a recommended study the appropriate condition-specific guideline.
  - There are significant potential adverse effects associated with the use of iodinated contrast media. These include hypersensitivity reactions, thyroid dysfunction, and contrast-induced nephropathy (CIN). Individuals with impaired renal function are at increased risk for CIN.<sup>2</sup>
  - Both contrast CT and MRI may be considered to have the same risk profile with renal failure (GFR <30 mL/min).
  - The use of CT contrast should proceed with caution in pregnant and breastfeeding individuals. There is a theoretical risk of contrast toxicity to the fetal and infant thyroid. The procedure can be performed if the specific need for that contrast-enhanced procedure outweighs risk to the fetus. Breastfeeding individuals may reduce this risk by choosing to pump and discard breast milk for 12-24 hours after the contrast injection.
  - CT without contrast may be appropriate if clinical criteria for CT with contrast are met AND the individual has:
    - Elevated blood urea nitrogen (BUN) and/or creatinine
    - Renal insufficiency
    - Allergies to iodinated contrast

- Thyroid disease which could be treated with I-131
- Diabetes
- Very elderly
- Urgent or emergent settings due to availability
- Trauma
- CT is superior to other imaging modalities in certain conditions including, but not limited to, the following:
  - Screening following trauma
  - Imaging pulmonary disease
  - Imaging abdominal and pelvic viscera
  - Imaging of complex fractures
  - Evaluation of inconclusive findings on Ultrasound or MRI, or if there is a contraindication to MRI
- More specific guidance for CT usage, including exceptions to this general guidance, can be found throughout the condition-specific guidelines.

### **Magnetic Resonance Imaging (MRI)**

- The AMA CPT<sup>®</sup> manual does not describe nor assign any minimum or maximum number of sequences for any MRI study. MRI protocols are often influenced by the individual's clinical situation and additional sequences are not uncommon. There are numerous MRI sequences that may be performed to evaluate specific clinical questions, and this technology is constantly undergoing development.
- Magnetic Resonance Imaging (MRI) utilizes the interaction between the intrinsic radiofrequency of certain molecules in the body (hydrogen in most cases) and a strong external magnetic field.
  - MRI is often superior for advanced imaging of soft tissues and can also define physiological processes in some instances (e.g., edema, loss of circulation [AVN], and increased vascularity [tumors]).
  - MRI does not use ionizing radiation and even non-contrast images have much higher soft tissue definition than CT or Ultrasound.
  - MRI typically takes much longer than either CT or Ultrasound, and for some individuals may require sedation. It is also much more sensitive to individual motion that can degrade image quality than either CT or Ultrasound.
- MRI Breast and MRI Chest are not interchangeable, as they focus detailed sequences on different adjacent body parts.
- MRI may be utilized either as the primary advanced imaging modality, or when further definition is needed based on CT or ultrasound imaging.
- Most orthopedic and dental implants are not magnetic. These include hip and knee replacements; plates, screws, and rods used to treat fractures; and cavity fillings. Yet,

all of these metal implants can distort the MRI image if near the part of the body being scanned.

- Other implants, however, may have contraindications to MRI. These include the following:
  - Pacemakers
  - ICD or heart valves
  - Metal implants in the brain
  - Metal implants in the eyes or ears
  - Infusion catheters and bullets or shrapnel
- CT can therefore be an alternative study to MRI in these scenarios.
- The contrast level and anatomic region in MRI imaging is specific to the clinical indication, as listed in the specific guideline sections.
- MRI utilizing Xenon Xe 129 for contrast is considered investigational and experimental at this time. MRI with or with and without contrast in these guidelines refers to MRI utilizing gadolinium for contrast.
- MRI is commonly performed without, without and with contrast.
  - Non-contrast imaging offers excellent tissue definition.
  - Imaging without and with contrast is commonly used when needed to better characterize tissue perfusion and vascularization.
    - Most contrast is gadolinium based and causes T2 brightening of the vascular and extracellular spaces.
    - Some specialized gadolinium and non-gadolinium contrast agents are available, and most commonly used for characterizing liver lesions.
  - MRI with contrast only is rarely appropriate and is usually used to better characterize findings on a recent inconclusive non-contrast MRI, commonly called a completion study.
  - MRI contrast is contraindicated in pregnant individuals.
  - More specific guidance for MRI contrast usage, including exceptions to this general guidance, can be found throughout the condition-specific guidelines.
- MRI may be preferred in individuals with renal failure and in individuals allergic to intravenous CT contrast.
  - Both contrast CT and MRI may be considered to have the same risk profile with renal failure (GFR <30 mL/min).<sup>2</sup>
  - Gadolinium can cause Nephrogenic Systemic Fibrosis (NSF). The greater the exposure to gadolinium in individuals with a low GFR (especially if on dialysis), the greater the chance of individuals developing NSF.
  - Multiple studies have demonstrated potential for gadolinium deposition following the use of gadolinium-based contrast agents (GBCAs) for MRI studies.<sup>3,4,5,6,7</sup> The U.S. Food and Drug Administration (FDA) has noted that there is currently no evidence to suggest that gadolinium retention in the brain is harmful and restricting

gadolinium-based contrast agents (GBCAs) use is not warranted at this time. It has been recommended that GBCA use should be limited to circumstances in which additional information provided by the contrast agent is necessary and the necessity of repetitive MRIs with GBCAs should be assessed.<sup>8</sup>

- A CT may be approved in place of an MRI when clinical criteria are met for MRI AND there is a contraindication to having an MRI (pacemaker, ICD, insulin pump, neurostimulator, etc.).
  - When replacing MRI with CT, contrast level matching should occur as follows:
    - MRI without contrast → CT without contrast
    - MRI without and with contrast → CT with contrast or CT without and with contrast
- The following situations may impact the appropriateness for MRI and or MR contrast:
  - Caution should be taken in the use of gadolinium in individuals with renal failure.
  - The use of gadolinium contrast agents is contraindicated during pregnancy unless the specific need for that procedure outweighs risk to the fetus.
  - MRI can be performed for non-ferromagnetic body metals (i.e., titanium), although some imaging facilities will consider it contraindicated if recent surgery, regardless of the metal type.
- MRI should not be used as a replacement for CT for the sole reason of avoidance of ionizing radiation when MRI is not supported in the condition-based guidelines, since it does not solve the problem of overutilization.
- MRI is superior to other imaging modalities in certain conditions including, but not limited to, the following:
  - Imaging the brain and spinal cord
  - Characterizing visceral and musculoskeletal soft tissue masses
  - Evaluating musculoskeletal soft tissues including ligaments and tendons
  - Evaluating inconclusive findings on ultrasound or CT
  - Individuals who are pregnant or have high radiation sensitivity
  - Suspicion, diagnosis, or surveillance of infections
- More specific guidance for MRI usage, including exceptions to this general guidance, can be found throughout the condition-specific guidelines.

### **Positron Emission Tomography (PET)**

- PET is a nuclear medicine study that uses a positron emitting radiotracer to create cross-sectional and volumetric images based on tissue metabolism.
- Conventional imaging (frequently CT, sometimes MRI or bone scan) of the affected area(s) drives much of initial and restaging and surveillance imaging for malignancy and other chronic conditions. PET is not indicated for surveillance imaging unless specifically stated in the condition-specific guideline sections.
- PET/MRI is generally not supported, see **PET-MRI (Preface-5.3)**.

- PET is rarely performed as a single modality, but is typically performed as a combined PET/CT.
  - The unbundling of PET/CT into separate PET and diagnostic CT CPT<sup>®</sup> codes is not supported, because PET/CT is done as a single study.
- PET/CT lacks the tissue definition of CT or MRI, but is fairly specific for metabolic activity based on the radiotracer used.
- Indications for PET/CT may include the following:
  - Oncologic Imaging for evaluation of tumor metabolic activity
  - Cardiac Imaging for evaluation of myocardial metabolic activity
  - Brain Imaging for evaluation of metabolic activity for procedural planning
- More specific guidance for PET usage, including exceptions to this general guidance, can be found throughout the condition-specific guidelines.

### **Overutilization of Advanced Imaging**

- A number of recent reports describe overutilization in many areas of advanced imaging and other procedures, which may include the following:
  - High-level testing without consideration of less invasive, lower cost options which may adequately address the clinical question at hand
  - Excessive radiation and costs with unnecessary testing
  - Defensive medical practice
  - CT without and with contrast (so called "double contrast studies") requests, which have few current indications
  - MRI requested in place of CT to avoid radiation without considering the primary indication for imaging
  - Adult CT settings and protocols used for smaller people and children
  - Unnecessary imaging procedures when the same or similar studies have already been conducted
- A review of the imaging or other relevant procedural histories of all individuals presenting for studies has been recognized as one of the more important processes that can be significantly improved. By recognizing that a duplicate or questionably indicated examination has been ordered for individuals, it may be possible to avoid exposing them to unnecessary risks.<sup>9,10</sup> To avoid these unnecessary risks, the precautions below should be considered:
  - The results of initial diagnostic tests or radiologic studies to narrow the differential diagnosis should be obtained prior to performing further tests or radiologic studies.
  - The clinical history should include a potential indication such as a known or suspected abnormality involving the body part for which the imaging study is being requested. These potential indications are addressed in greater detail within the applicable guidelines.

- The results of the requested imaging procedures should be expected to have an impact on individual management or treatment decisions.
- Repeat imaging studies are not generally necessary unless there is evidence of disease progression, recurrence of disease, and/or the repeat imaging will affect an individual's clinical management.
- Pre-operative imaging/pre-surgical planning imaging/pre-procedure imaging is not indicated if the surgery/procedure is not indicated. Once the procedure has been approved or if the procedure does not require prior authorization, the appropriate pre-procedural imaging may be approved.



## References (Preface-3)

v2.0.2024

1. Bettmann MA. Frequently Asked Questions: Iodinated Contrast Agents. *RadioGraphics*. 2004;24(suppl\_1):S3-S10. doi: 10.1148/rg.24si045519.
2. Andreucci M, Solomon R, Tasanarong A. Side Effects of Radiographic Contrast Media: Pathogenesis, Risk Factors, and Prevention. *BioMed Res Int*. 2014;2014:1-20. doi: 10.1155/2014/741018.
3. McDonald RJ, McDonald JS, Kallmes DF, et al. Intracranial Gadolinium Deposition after Contrast-enhanced MR Imaging. *Radiology*. 2015;275(3):772-782. doi: 10.1148/radiol.15150025.
4. Kanda T, Ishii K, Kawaguchi H, Kitajima K, Takenaka D. High Signal Intensity in the Dentate Nucleus and Globus Pallidus on Unenhanced T1-weighted MR Images: Relationship with Increasing Cumulative Dose of a Gadolinium-based Contrast Material. *Radiology*. 2014;270(3):834-841. doi: 10.1148/radiol.13131669.
5. Olchowcy C, Cebulski K, Łasecki M, et al. The presence of the gadolinium-based contrast agent depositions in the brain and symptoms of gadolinium neurotoxicity - A systematic review. Mohapatra S, ed. *PLOS ONE*. 2017;12(2):e0171704. doi: 10.1371/journal.pone.0171704.
6. Ramalho J, Castillo M, AlObaidy M, et al. High Signal Intensity in Globus Pallidus and Dentate Nucleus on Unenhanced T1-weighted MR Images: Evaluation of Two Linear Gadolinium-based Contrast Agents. *Radiology*. 2015;276(3):836-844. doi:10.1148/radiol.2015150872.
7. Radbruch A, Weberling LD, Kieslich PJ, et al. Intraindividual Analysis of Signal Intensity Changes in the Dentate Nucleus After Consecutive Serial Applications of Linear and Macrocyclic Gadolinium-Based Contrast Agents. *Invest Radiol*. 2016;51(11):683-690. doi: 10.1097/rni.0000000000000308.
8. FDA Warns That Gadolinium-Based Contrast Agents (GBCAs) Are Retained in the Body; Requires New Class Warnings. <https://www.fda.gov/media/109825/download>.
9. Amis ES, Butler PF, Applegate KE, et al. American College of Radiology White Paper on Radiation Dose in Medicine. *J Am Coll Radiol*. 2007;4(5):272-284. doi: 10.1016/j.jacr.2007.03.002.
10. Powell AC, Long JW, Kren EM, Gupta AK, Levin DC. Evaluation of a Program for Improving Advanced Imaging Interpretation. *J Patient Saf*. 2019;15(1):69-75. doi: 10.1097/PTS.000000000000034.5.
11. FDA. White Paper: Initiative to Reduce Unnecessary Radiation Exposure from Medical Imaging. Page Last Updated: 06/14/2019. <https://www.fda.gov/Radiation-EmittingProducts/RadiationSafety/RadiationDoseReduction/ucm199994.htm>.
12. Update on FDA approach to safety issue of gadolinium retention after administration of gadolinium-based contrast agents. <https://www.fda.gov/media/116492/download>.
13. Blumfield E, Swenson DW, Iyer RS, Stanescu AL. Gadolinium-based contrast agents — review of recent literature on magnetic resonance imaging signal intensity changes and tissue deposits, with emphasis on pediatric patients. *Pediatr Radiol*. 2019;49(4):448-457. doi: 10.1007/s00247-018-4304-8.
14. American College of Radiology. ACR – SPR – SRU Practice Parameter for the Performing and Interpreting Diagnostic Ultrasound Examinations. Revised 2017. (Resolution 32). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-Perf-Interpret.pdf>.
15. American College of Radiology. ACR–SPR Practice Parameter for Performing FDG-PET/CT in Oncology. Revised 2021. (Resolution 20). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/FDG-PET-CT.pdf>.
16. American College of Radiology. ACR Practice Parameter for Performing and Interpreting Magnetic Resonance Imaging (MRI). Revised 2017. (Resolution 10). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Perf-Interpret.pdf>.
17. American College of Radiology. ACR Practice Parameter for Performing and Interpreting Diagnostic Computed Tomography (CT). Revised 2017. (Resolution 22). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CT-Perf-Interpret.pdf>.
18. Lohrke J, Frenzel T, Endrikat J, et al. 25 Years of Contrast-Enhanced MRI: Developments, Current Challenges and Future Perspectives. *Adv Ther*. 2016;33(1):1-28. doi: 10.1007/s12325-015-0275-4.
19. Implementation Guide: Medicaid State Plan Eligibility Eligibility Groups Mandatory Coverage Infants and Children under Age 19. Available at: <https://www.hhs.gov/guidance/document/implementation-guide-medicaid-state-plan-eligibility-eligibility-groups-aeu-mandatory-2>.

20. History and Physicals - Understanding the Requirements. Available at: <https://www.jointcommission.org/standards/standard-faqs/hospital-and-hospital-clinics/provision-of-care-treatment-and-services-pc/000002272/>.
21. Mammarrappallil JG, Rankine L, Wild JM, Driehuys B. New Developments in Imaging Idiopathic Pulmonary Fibrosis With Hyperpolarized Xenon Magnetic Resonance Imaging. *J Thorac Imaging*. 2019;34(2):136-150. doi: 10.1097/rti.0000000000000392.
22. Wang JM, Robertson SH, Wang Z, et al. Using hyperpolarized <sup>129</sup>Xe MRI to quantify regional gas transfer in idiopathic pulmonary fibrosis. *Thorax*. 2017;73(1):21-28. doi: 10.1136/thoraxjnl-2017-210070.



# Coding Issues (Preface-4)

---

## Guideline

---

3D Rendering (Preface-4.1)

CT-, MR-, or Ultrasound-Guided Procedures (Preface-4.2)

Unlisted Procedures/Therapy Treatment Planning (Preface-4.3)

CPT<sup>®</sup> 76380 Limited or Follow-up CT (Preface-4.5)

SPECT/CT Imaging (Preface-4.6)

CPT<sup>®</sup> 76140 Interpretation of an Outside Study (Preface-4.7)

Quantitative MR Analysis of Tissue Composition (Preface-4.8)

HCPCS Codes (Preface-4.9)

References (Preface-4)

## 3D Rendering (Preface-4.1)

PRF.CD.0004.1.UOH

v2.0.2024

### CPT<sup>®</sup> 76376 and CPT<sup>®</sup> 76377

- Both codes require concurrent supervision of the image post-processing 3D manipulation of the volumetric data set and image rendering.
  - Concurrent supervision is defined as active physician participation in and monitoring of the reconstruction process including design of the anatomic region that is to be reconstructed; determination of the tissue types and actual structures to be displayed (e.g., bone, organs, and vessels); determination of the images or cine loops that are to be archived; and, monitoring and adjustment of the 3D work product. The American College of Radiology (ACR) recommends that it is best to document the physician's supervision or participation in the 3D reconstruction of images.
- These two codes differ in the need for and use of an independent workstation for post-processing.
  - CPT<sup>®</sup> 76376 reports procedures not requiring image post-processing on an independent workstation.
  - CPT<sup>®</sup> 76377 reports procedures that require image post-processing on an independent workstation.
- These 3D rendering codes should not be used for 2D reformatting.
- Two-dimensional reconstruction (e.g., reformatting an axial scan into the coronal plane) is now included in all cross-sectional imaging base codes and is not separately reimbursable.
- The codes used to report 3D rendering for ultrasound and echocardiography are also used to report the 3D post processing work on CT, MRI, and other tomographic modalities.
- Providers may be required to obtain prior authorization on these 3D codes even if prior authorization is not required for the echocardiography and/or ultrasound procedure codes. It may appear that UnitedHealthcare pre-authorizes echocardiography and/or ultrasound when, in fact, it may only be the 3D code that needs the prior authorization.
- CPT<sup>®</sup> codes for 3D rendering should not be billed in conjunction with computer-aided detection (CAD), MRA, CTA, nuclear medicine SPECT studies, PET, PET/CT, Mammogram, MRI Breast, US Breast, CT Colonography (virtual colonoscopy), Cardiac MRI, Cardiac CT, or Coronary CTA studies.

- CPT<sup>®</sup> 76377 (3D rendering requiring image post-processing on an independent workstation) or CPT<sup>®</sup> 76376 (3D rendering not requiring image post-processing on an independent workstation) can be considered in the following clinical scenarios:
  - Bony conditions:
    - Evaluation of congenital skull abnormalities in newborns, infants, and toddlers (usually for pre-operative planning)
    - Complex fractures (comminuted or displaced)/dislocations of any joint (for pre-operative planning when conventional imaging is insufficient)
    - Spine fractures, pelvic/acetabulum fractures, intra-articular fractures (for pre-operative planning when conventional imaging is insufficient)
    - Pre-operative planning for other complex surgical cases
    - Complex facial fractures
  - Pre-operative planning for other complex surgical cases
  - Cerebral angiography
  - Pelvis conditions:
    - Uterine intra-cavitary lesion when initial US is equivocal: See **Abnormal Uterine Bleeding (AUB) (PV-2.1)** and **Leiomyoma/Uterine Fibroids (PV-12.1)** in the Pelvis Imaging Guidelines.
    - Hydrosalpinxes or peritoneal cysts when initial US is indeterminate: See **Complex Adnexal Masses (PV-5.3)** in the Pelvis Imaging Guidelines.
    - Lost IUD (inability to feel or see IUD string) with initial US: See **Intrauterine Device (PV-10.1)** in the Pelvis Imaging Guidelines.
    - Uterine anomalies with initial US: See **Uterine Anomalies (PV-14.1)** in the Pelvis Imaging Guidelines.
    - Infertility: See **Initial Infertility Evaluation, Female (PV-9.1)** in the Pelvis Imaging Guidelines.
  - Abdomen conditions:
    - CT Urogram: See **Hematuria and Hydronephrosis (AB-39)** in the Abdomen Imaging Guidelines.
    - MRCP: See **MR Cholangiopancreatography (MRCP) (AB-27)** in the Abdomen Imaging Guidelines.

## CT-, MR-, or Ultrasound-Guided Procedures (Preface-4.2)

PRF.CD.0004.2.A

v2.0.2024

- CT-, MR-, and Ultrasound-guidance procedure codes contain all of the imaging necessary to guide a needle or catheter. It is inappropriate to routinely bill a diagnostic procedure code in conjunction with a guidance procedure code.
- Imaging studies performed as part of a CT-, MR-, or Ultrasound-guided procedure should be reported using the CPT<sup>®</sup> codes in the following table:

**TABLE: Imaging Guidance Procedure Codes**

CPT <sup>®</sup>	Description
<b>19085</b>	Biopsy, breast, with placement of breast localization device(s), when performed, and imaging of the biopsy specimen, when performed, percutaneous; first lesion, including MR guidance
<b>19086</b>	Biopsy, breast, with placement of breast localization device(s), when performed, and imaging of the biopsy specimen, when performed, percutaneous; each additional lesion, including MR guidance
<b>75989</b>	Imaging guidance for percutaneous drainage with placement of catheter (all modalities)
<b>76942</b>	Ultrasonic guidance for needle placement
<b>77011</b>	CT guidance for stereotactic localization
<b>77012</b>	CT guidance for needle placement
<b>77013</b>	CT guidance for, and monitoring of parenchymal tissue ablation
<b>77021</b>	MR guidance for needle placement
<b>77022</b>	MR guidance for, and monitoring of parenchymal tissue ablation

### **CPT® 19085 and CPT® 19086**

- The proper way to bill an MRI-guided breast biopsy is CPT® 19085 (Biopsy, breast, with placement of breast localization device(s), when performed, and imaging of the biopsy specimen, when performed, percutaneous; first lesion, including MR guidance). Additional lesions should be billed using CPT® 19086.
  - **CPT® 77021** (MR guidance for needle placement) is not an appropriate code for a breast biopsy.

### **CPT® 75989**

- This code is used to report imaging guidance for a percutaneous drainage procedure in which a catheter is left in place.
- This code can be used to report whether the drainage catheter is placed under fluoroscopy, Ultrasound-, CT-, or MR-guidance modality.

### **CPT® 77011**

- A stereotactic CT localization scan is frequently obtained prior to sinus surgery. The dataset is then loaded into the navigational workstation in the operating room for use during the surgical procedure. The information provides exact positioning of surgical instruments with regard to the individual's 3D CT images.<sup>3</sup>
- In most cases, the pre-operative CT is a technical-only service that does not require interpretation by a radiologist.
  - The imaging facility should report CPT® 77011 when performing a scan not requiring interpretation by a radiologist.
  - If a diagnostic scan is performed and interpreted by a radiologist, the appropriate diagnostic CT code (e.g., CPT® 70486) should be used.
  - It is not appropriate to report both CPT® 70486 and CPT® 77011 for the same CT stereotactic localization imaging session.
  - 3D Rendering (CPT® 76376 or CPT® 76377) should not be reported in conjunction with CPT® 77011 (or CPT® 70486 if used). The procedure inherently generates a 3D dataset.

### **CPT® 77012 (CT) and CPT® 77021 (MR)**

- These codes are used to report imaging guidance for needle placement during biopsy, aspiration, and other percutaneous procedures.
- They represent the radiological supervision and interpretation of the procedure and are often billed in conjunction with surgical procedure codes.
  - For example, CPT® 77012 is reported when CT guidance is used to place the needle for a conventional arthrogram.
  - Only codes representing percutaneous surgical procedures should be billed with CPT® 77012 and CPT® 77021. It is inappropriate to use with surgical codes for open, excisional, or incisional procedures.

- **CPT<sup>®</sup> 77021** (MR guidance for needle placement) is not an appropriate code for breast biopsy.
  - CPT<sup>®</sup> 19085 would be appropriate for the first breast biopsy site and CPT<sup>®</sup> 19086 would be appropriate for additional concurrent biopsies.

**CPT<sup>®</sup> 77013 (CT) and CPT<sup>®</sup> 77022 (MR)**

- These codes include the initial guidance to direct a needle electrode to the tumor(s), monitoring for needle electrode repositioning within the lesion, and as necessary for multiple ablations to coagulate the lesion and confirmation of satisfactory coagulative necrosis of the lesion(s) and comparison to pre-ablation images.
  - **NOTE:** CPT<sup>®</sup> 77013 should only be used for non-bone ablation procedures.
    - CPT<sup>®</sup> 20982 includes CT guidance for bone tumor ablations.
  - Only codes representing percutaneous surgical procedures should be billed with CPT<sup>®</sup> 77013 and CPT<sup>®</sup> 77022. It is inappropriate to use with surgical codes for open, excisional, or incisional procedures.
- CPT<sup>®</sup> 77012 and CPT<sup>®</sup> 77021 (as well as guidance codes CPT<sup>®</sup> 76942 [US], and CPT<sup>®</sup> 77002 - CPT<sup>®</sup> 77003 [fluoroscopy]) describe radiologic guidance by different modalities.
  - Only one unit of any of these codes should be reported per individual encounter (date of service). The unit of service is considered to be the individual encounter, not the number of lesions, aspirations, biopsies, injections, or localizations.

# Unlisted Procedures/Therapy Treatment Planning (Preface-4.3)

PRF.CD.0004.3.UOH

v2.0.2024

CPT <sup>®</sup>	Description
76497	Unlisted CT procedure (e.g., diagnostic or interventional)
76498	Unlisted MR procedure (e.g., diagnostic or interventional)
78999	Unlisted procedure, diagnostic nuclear medicine

- These unlisted codes should be reported whenever a diagnostic or interventional CT or MR study is performed in which an appropriate anatomic site-specific code is not available.
  - A Category III code that describes the procedure performed must be reported rather than an unlisted code if one is available.
- CPT<sup>®</sup> 76497 or CPT<sup>®</sup> 76498 (Unlisted CT or MRI procedure) can be considered in the following clinical scenarios:
  - Studies done for navigation and planning for neurosurgical procedures (i.e., Stealth or Brain Lab Imaging)<sup>1,2</sup>
  - Custom joint arthroplasty planning (not as an alternative recommendation): See **Osteoarthritis (MS-12.1)** in the Musculoskeletal Imaging Guidelines.
  - Any procedure/surgical planning if thinner cuts or different positional acquisition (than those on the completed diagnostic study) are needed. These could include navigational bronchoscopy: See **Navigational Bronchoscopy (CH-1.7)** in the Chest Imaging Guidelines.

## Therapy Treatment Planning

- Radiation Therapy Treatment Planning: See **Unlisted Procedure Codes in Oncology (ONC-1.5)** in the Oncology Imaging Guidelines.

# CPT<sup>®</sup> 76380 Limited or Follow-up CT (Preface-4.5)

PRF.CD.0004.5.UOH

v2.0.2024

- CPT<sup>®</sup> 76380 describes a limited or follow-up CT scan. The code is used to report any CT scan, for any given area of the body, in which the work of a full diagnostic code is not performed.
- Common examples include, but are not limited to, the following:
  - Limited sinus CT imaging protocol
  - Limited or follow-up slices through a known pulmonary nodule
  - Limited slices to assess a non-healing fracture (such as the clavicle)
- Limited CT (CPT<sup>®</sup> 76380) is not indicated for treatment planning purposes. See **Unlisted Procedure Codes in Oncology (ONC-1.5)** in the Oncology Imaging Guidelines.
- It is inappropriate to report CPT<sup>®</sup> 76380, in conjunction with other diagnostic CT codes, to cover 'extra slices' in certain imaging protocols.
  - There is no specific number of sequences or slices defined in any CT CPT<sup>®</sup> code definition.
  - The AMA, in *CPT<sup>®</sup> 2019*, does not describe nor assign any minimum or maximum number of sequences or slices for any CT study.
    - A few additional slices or sequences are not uncommon.
    - CT imaging protocols are often influenced by the individual's clinical situation. Sometimes the protocols require more time and sometimes less.



## **SPECT/CT Imaging (Preface-4.6)**

**PRF.CD.0004.6.A**

**v2.0.2024**

- SPECT/CT involves SPECT (Single Photon Emission Computed Tomography) nuclear medicine imaging and CT for optimizing location, accuracy, and attenuation correction and combines functional and anatomic information.
  - Common studies using this modality include  $^{123}\text{I}$ - or  $^{131}\text{I}$ -Metaiodobenzylguanidine (MIBG) and octreotide scintigraphy for neuroendocrine tumors.
- Hybrid Nuclear/CT scan can be reported as CPT<sup>®</sup> 78830 (single area and single day), CPT<sup>®</sup> 78831 (2 or more days), or CPT<sup>®</sup> 78832 (2 areas with one day and 2-day study).
- CPT<sup>®</sup> 78072 became effective January 1, 2013 for SPECT/CT parathyroid nuclear imaging.

# CPT<sup>®</sup> 76140 Interpretation of an Outside Study (Preface-4.7)

PRF.CD.0004.7.UOH

v2.0.2024

- It is inappropriate to use diagnostic imaging codes for interpretation of a previously performed exam that was completed at another facility.
  - If the outside exam is being used for comparison with a current exam, the diagnostic code for the current examination includes comparison to the prior study.<sup>4</sup>
  - CPT<sup>®</sup> 76140 is the appropriate code to use for an exam which was completed elsewhere and a secondary interpretation of the images is requested.<sup>5</sup>

# Quantitative MR Analysis of Tissue Composition (Preface-4.8)

PRF.CD.0004.8.A

v2.0.2024

- Category III CPT<sup>®</sup> codes for quantitative analysis of multiparametric-MR (mp-MRI) data with and without an associated diagnostic MRI have been established. Quantitative mp-MRI uses software to analyze tissue physiology of visceral organs and other anatomic structures non-invasively. At present, these procedures are primarily being used in clinical trials and there is no widely recommended indications in clinical practice. As such, these procedures are considered to be investigational and experimental for coverage purposes.
  - CPT<sup>®</sup> 0648T (without diagnostic MRI) and CPT<sup>®</sup> 0649T (with diagnostic MRI) refer to data analysis with and without associate imaging of a single organ, with its most common use being LiverMultiScan (LMS).
    - See **Fatty Liver (AB-29.2)** in the Abdomen Imaging Guidelines.
  - CPT<sup>®</sup> 0697T (without diagnostic MRI) and CPT<sup>®</sup> 0698T (with diagnostic MRI) refer to data analysis with and without associate imaging of a multiple organs, with its most common use being CoverScan.

## HCPCS Codes (Preface-4.9)

PRF.CD.0004.9.UOH

v2.0.2024

- Healthcare Common Procedure Coding System (HCPCS) codes are utilized by some hospitals in favor of the typical Level-III CPT<sup>®</sup> codes. These codes are typically 4 digits preceded by a C or S.<sup>6</sup>
  - Many of these codes have similar code descriptions to Level-III CPT<sup>®</sup> codes (i.e., C8931 – MRA with dye, Spinal Canal; and, CPT<sup>®</sup> 72159 – MRA Spinal Canal).
  - If cases are submitted with HCPCS codes with similar code descriptions to the typical Level-III CPT<sup>®</sup> codes, those procedures should be managed in the same manner as the typical CPT<sup>®</sup> codes.
  - HCPCS code management is discussed further in the applicable guideline sections.
- Requests for many Healthcare Common Procedure Coding System (HCPCS) codes, including non-specific codes such as S8042 (Magnetic resonance imaging [MRI], low-field), should be redirected to a more appropriate and specific CPT<sup>®</sup> code. Exceptions are noted in the applicable guideline sections.

## References (Preface-4)

**v2.0.2024**

1. Society of Nuclear Medicine and Molecular Imaging Coding Corner. Available at: <http://www.snmmi.org/ClinicalPractice/CodingCornerPT.aspx?ItemNumber=1786>.
2. Intraoperative MR. Brainlab. Available at: <https://www.brainlab.com/surgery-products/overview-neurosurgery-products/intraoperative-mr/>
3. Experience the Advanced 3D Sinus Surgery Planning with Scopis Building Blocks planning software. Scopis Planning. Available at: <http://planning.scopis.com/>.
4. ACR Radiology Coding Source™ March-April 2007 Q and A. Available at: <https://www.acr.org/Advocacy-and-Economics/Coding-Source/ACR-Radiology-Coding-Source-March-April-2007-Q-and-A>.
5. Chung CY, Alson MD, Duszak R, Degnan AJ. From imaging to reimbursement: what the pediatric radiologist needs to know about health care payers, documentation, coding and billing. *Pediatr Radiol*. 2018;48(7):904-914. doi: 10.1007/s00247-018-4104-1.
6. HCPCS - General Information from CMS.gov. Available at: [www.cms.gov/medicare/coding/medhcpcsgeninfo](http://www.cms.gov/medicare/coding/medhcpcsgeninfo).

# Whole-Body Imaging (Preface-5)

---

## Guideline

---

Whole-Body CT Imaging (Preface-5.1)  
Whole-Body MR Imaging (Preface-5.2)  
PET-MRI (Preface-5.3)  
References (Preface-5)

## Whole-Body CT Imaging (Preface-5.1)

PRF.WB.0005.1.UOH

v2.0.2024

- Whole-body CT or LifeScan (CT Brain, Chest, Abdomen, and Pelvis) for screening of asymptomatic individuals is not indicated. The performance of whole-body screening CT examinations in healthy individuals does not meet any of the current validity criteria for screening studies and there is no clear documentation of benefit versus radiation risk.
- Whole-body low-dose CT is supported for oncologic staging in Multiple Myeloma. See **Multiple Myeloma and Plasmacytomas (ONC-25)** in the Oncology Imaging Guidelines.

## Whole-Body MR Imaging (Preface-5.2)

PRF.WB.0005.2.A

v2.0.2024

- Whole-body MRI (WBMRI) is, with the exception of select cancer predisposition syndromes and autoimmune conditions discussed below, generally not supported at this time due to lack of standardization in imaging technique and lack of evidence that WBMRI improves outcome for any individual disease state.
  - While WBMRI has the benefit of whole-body imaging and lack of radiation exposure, substantial variation still exists in the number of images, type of sequences (STIR vs. diffusion weighting, for example), and contrast agent(s) used.
- Coding considerations:
  - There are no established CPT<sup>®</sup> or HCPCS codes for reporting WBMRI.
  - WBMRI is at present only reportable using CPT<sup>®</sup> 76498. All other methods of reporting whole-body MRI are inappropriate including the following:
    - Separate diagnostic MRI codes for multiple individual body parts
    - MRI Bone Marrow Supply (CPT<sup>®</sup> 77084)
- Disease-specific considerations:
  - Cancer screening:
    - Interval WBMRI is recommended for cancer screening in individuals with select cancer predisposition syndromes. Otherwise, WBMRI has not been shown to improve outcomes for cancer screening.
      - For additional information, see **Li-Fraumeni Syndrome (LFS) (PEDONC-2.2)** , **Hereditary Paraganglioma-Pheochromocytoma (HPP) Syndromes (PEDONC-2.13)** , or **Constitutional Mismatch Repair Deficiency (CMMRD or Turcot Syndrome) (PEDONC-2.15)** in the Pediatric Oncology Imaging Guidelines.
  - Cancer staging and restaging:
    - While the feasibility of WBMRI has been established, data remain conflicting on whether WBMRI is of equivalent diagnostic accuracy compared with standard imaging modalities such as CT, scintigraphy, and PET imaging.
    - Evidence has not been published establishing WBMRI as a standard evaluation for any type of cancer.
  - Autoimmune disease:
    - WBMRI can be approved in some situations for individuals with chronic recurrent multifocal osteomyelitis.
      - For additional information, see **Chronic Recurrent Multifocal Osteomyelitis (PEDMS-10.2)** in the Pediatric Musculoskeletal Imaging Guidelines.



## PET-MRI (Preface-5.3)

PRF.WB.0005.3.A

v2.0.2024

- PET-MRI is generally not supported for a vast majority of oncologic and neurologic conditions due to lack of standardization in imaging technique and interpretation. However, it may be appropriate in select circumstances when the following criteria are met:
  - The individual meets condition-specific guidelines for PET-MRI OR
  - The individual meets ALL of the following:
    - The individual is a pediatric patient or being treated under a pediatric guideline and treatment plan AND
    - The individual meets guideline criteria for PET-CT, **AND**
    - PET-CT is not available at the treating institution, **AND**
    - The provider requests PET-MRI in lieu of PET-CT
- When the above criteria are met, PET-MRI may be reported using the code combination of PET Whole-Body (CPT<sup>®</sup> 78813) and MRI Unlisted (CPT<sup>®</sup> 76498). All other methods of reporting PET-MRI are inappropriate.
  - When clinically appropriate, diagnostic MRI codes may be indicated at the same time as the PET-MRI code combination.
- For more information, see **PET Imaging in Pediatric Oncology (PEDONC-1.4)** in the Pediatric Oncology Imaging Guidelines, and **PET Brain Imaging (PEDHD-2.3)** and **Special Imaging Studies in Evaluation for Epilepsy Surgery (PEDHD-6.3)** in the Pediatric Head Imaging Guidelines.

## References (Preface-5)

**v2.0.2024**

1. Villani A, Tabori U, Schiffman J, et al. Biochemical and imaging surveillance in germline TP53 mutation carriers with Li-Fraumeni syndrome: a prospective observational study. *Lancet Oncol.* 2011;12(6):559-567. doi: 10.1016/S1470-2045(11)70119-X.
2. Siegel MJ, Acharyya S, Hoffer FA, et al. Whole-Body MR Imaging for Staging of Malignant Tumors in Pediatric Patients: Results of the American College of Radiology Imaging Network 6660 Trial. *Radiology.* 2013;266(2):599-609. doi: 10.1148/radiol.12112531.
3. Antoch G. Whole-Body Dual-Modality PET/CT and Whole-Body MRI for Tumor Staging in Oncology. *JAMA.* 2003;290(24):3199. doi: 10.1001/jama.290.24.3199.
4. Lauenstein TC, Semelka RC. Emerging techniques: Whole-body screening and staging with MRI. *J Magn Reson Imaging.* 2006;24(3):489-498. doi: 10.1002/jmri.20666.
5. Khanna G, Sato TSP, Ferguson P. Imaging of Chronic Recurrent Multifocal Osteomyelitis. *RadioGraphics.* 2009;29(4):1159-1177. doi: 10.1148/rg.294085244.
6. Ferguson PJ, Sandu M. Current Understanding of the Pathogenesis and Management of Chronic Recurrent Multifocal Osteomyelitis. *Curr Rheumatol Rep.* 2012;14(2):130-141. doi: 10.1007/s11926-012-0239-5.
7. National Comprehensive Cancer Network<sup>®</sup> (NCCN<sup>®</sup>). NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines<sup>®</sup>): Genetic/Familial High Risk Assessment: Breast, Ovarian, and Pancreatic. Version 3.2023. February 13, 2023. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines<sup>®</sup>) for Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic V.3.2023. ©National Comprehensive Cancer Network, Inc. 2023. All rights reserved. Accessed July 10, 2023. The NCCN Guidelines<sup>®</sup> and illustrations herein may not be reproduced in any form for any purpose without the express written permission of the NCCN. To view the most recent and complete version of the NCCN Guidelines<sup>®</sup>, go online to NCCN.org.

# References (Preface-6)

---

## Guideline

---

### References (Preface-6.1)

## References (Preface-6.1)

---

**PRF.RF.0006.1.A**

**v2.0.2024**

- Complete reference citations for the journal articles are embedded within the body of the guidelines and/or may be found on the Reference pages at the end of some guideline sections.
- The website addresses for certain references are included in the body of the guidelines but are not hyperlinked to the actual website.
- The website address for the American College of Radiology (ACR) Appropriateness Criteria<sup>®</sup> is <http://www.acr.org>.

# Copyright Information (Preface-7)

---

## Guideline

---

Copyright Information (Preface-7.1)

## Copyright Information (Preface-7.1)

PRF.CI.0007.1.UOH

v2.0.2024

- ©2024 United HealthCare Services, Inc. All rights reserved. No part of these materials may be changed, reproduced, or transmitted in any form or by any means, electronic or mechanical, including photocopying or recording, or in any information storage or retrieval system, without the prior express written permission of United HealthCare Services, Inc.

# Trademarks (Preface-8)

---

## Guideline

---

Trademarks (Preface-8.1)

## Trademarks (Preface-8.1)

PRF.TM.0008.1.A

v2.0.2024

- **CPT® (Current Procedural Terminology)** is a registered trademark of the American Medical Association (AMA). **CPT®** five-digit codes, nomenclature, and other data are copyright 2024 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values, or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.



# General Guidelines (MS-1)

---

## Guideline

---

Procedure Codes Associated with Musculoskeletal Imaging (MS)  
General Guidelines (MS-1.0)  
References (MS-1)

# Procedure Codes Associated with Musculoskeletal Imaging (MS)

MS.GG.ProcedureCodes.A

v1.0.2024

MRI/MRA	CPT®
MRI Upper Extremity, other than joint, without contrast	73218
MRI Upper Extremity, other than joint, with contrast	73219
MRI Upper Extremity, other than joint, without and with contrast	73220
MRI Upper Extremity, any joint, without contrast	73221
MRI Upper Extremity, any joint, with contrast	73222
MRI Upper Extremity, any joint, without and with contrast	73223
MR Angiography Upper Extremity without or with contrast	73225
MRI Lower Extremity, other than joint, without contrast	73718
MRI Lower Extremity, other than joint, with contrast	73719
MRI Lower Extremity, other than joint, without and with contrast	73720
MRI Lower Extremity, any joint, without contrast	73721
MRI Lower Extremity, any joint, with contrast	73722
MRI Lower Extremity, any joint, without and with contrast	73723
MR Angiography Lower Extremity without or with contrast	73725
MRI Pelvis without contrast	72195
MRI Pelvis with contrast	72196

<b>MRI/MRA</b>	<b>CPT®</b>
MRI Pelvis without and with contrast	72197

<b>CT/CTA</b>	<b>CPT®</b>
CT Upper Extremity without contrast	73200
CT Upper Extremity with contrast	73201
CT Upper Extremity without and with contrast	73202
CT Angiography Upper Extremity without and with contrast	73206
CT Lower Extremity without contrast	73700
CT Lower Extremity with contrast	73701
CT Lower Extremity without and with contrast	73702
CT Angiography Lower Extremity without and with contrast	73706
CT Pelvis without contrast	72192
CT Pelvis with contrast	72193
CT Pelvis without and with contrast	72194
Bone Mineral Density CT, one or more sites, axial skeleton	77078

<b>Ultrasound</b>	<b>CPT®</b>
Ultrasound, complete joint (ie, joint space and peri-articular soft tissue structures) real-time with image documentation	76881
Ultrasound, limited, joint or other nonvascular extremity structure(s) (e.g., joint space, peri-articular tendon[s], muscle[s], nerve[s], other soft tissue structure[s], or soft tissue mass[es]), real-time with image documentation	76882

<b>Ultrasound</b>	<b>CPT®</b>
Ultrasound, pelvic (nonobstetric), real time with image documentation	76857

<b>Nuclear Medicine</b>	<b>CPT®</b>
Bone Marrow Imaging, Limited	78102
Bone Marrow Imaging, Multiple	78103
Bone Marrow Imaging, Whole Body	78104
Bone or Joint Imaging Limited	78300
Bone or Joint Imaging Multiple	78305
Bone Scan Whole Body	78306
Bone Scan 3 Phase Study	78315
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, single area (e.g., head, neck, chest, pelvis), single day imaging	78800
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, 2 or more areas (eg, abdomen and pelvis, head and chest), 1 or more days imaging or single area imaging over 2 or more days	78801
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, whole body, single day imaging	78802
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT), single area (e.g., head, neck, chest, pelvis), single day imaging	78803

<b>Nuclear Medicine</b>	<b>CPT<sup>®</sup></b>
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT) with concurrently acquired computed tomography (CT) transmission scan for anatomical review, localization and determination/detection of pathology, single area (e.g., head, neck, chest, pelvis), single day imaging	78830
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT), minimum 2 areas (e.g., pelvis and knees, abdomen and pelvis), single day imaging, or single area imaging over 2 or more days	78831
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT) with concurrently acquired computed tomography (CT) transmission scan for anatomical review, localization and determination/detection of pathology, minimum 2 areas (e.g., pelvis and knees, abdomen and pelvis), single day imaging, or single area imaging over 2 or more days	78832

## General Guidelines (MS-1.0)

MS.GG.0001.0.A

v1.0.2024

- Before advanced diagnostic imaging can be considered, there must be an in-person clinical evaluation as well as a clinical re-evaluation after a trial of failed conservative treatment; the clinical re-evaluation may consist of an in-person evaluation or other meaningful contact with the provider's office such as email, web or telephone communications.
- An in-person clinical evaluation for the current episode of the condition is required to have been performed before advanced imaging can be considered. This may have been either the initial clinical evaluation or the clinical re-evaluation.
- The in-person clinical evaluation should include a relevant history and physical examination, appropriate laboratory studies, and non-advanced imaging modalities. Other forms of meaningful contact (e.g., telephone call, electronic mail, telemedicine, or messaging) are not acceptable as an in-person evaluation.
- Prior to advanced imaging consideration, the results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider of the advanced imaging study for all musculoskeletal conditions, unless otherwise noted in the guidelines.
  - Initial plain x-ray can rule out those situations that do not often require advanced imaging, such as osteoarthritis, acute/healing fracture, dislocation, osteomyelitis, acquired/congenital deformities, and tumors of bone amenable to biopsy or radiation therapy (in known metastatic disease), etc.
  - X-ray may provide complementary clinical information regarding detailed bony anatomy, and may assist with preoperative planning when surgery is being contemplated.
  - X-ray may provide clinically significant details for soft tissue masses, such as soft tissue calcification, presence or absence of phleboliths, radiographic density, and effect on adjacent bone.
  - X-ray often has a larger field of view than MRI or CT and has the potential to identify more proximal or distal pathology in an extremity.
- Clinical re-evaluation is required prior to consideration of advanced diagnostic imaging to document failure of significant clinical improvement following a recent (within 3 months) six week trial of provider-directed conservative treatment. Clinical re-evaluation can include documentation of an in-person encounter or documentation of other meaningful contact with the requesting provider's office by the individual (e.g. telephone call, electronic mail, telemedicine, or messaging).
- Provider-directed conservative treatment may include rest, ice, compression, and elevation (R.I.C.E.), non-steroidal anti-inflammatories (NSAIDs), narcotic

and non-narcotic analgesic medications, oral or injectable corticosteroids, viscosupplementation injections, a provider-directed home exercise program, cross-training, and/or physical/occupational therapy or immobilization by splinting/casting/bracing.

- Orthopedic specialist evaluation can be helpful in determining the need for advanced imaging.
  - The need for repeat advanced imaging should be carefully considered and may not be indicated if prior imaging has been performed.
  - Serial advanced imaging, whether CT or MRI, for surveillance of healing or recovery from musculoskeletal disease is not supported by the medical evidence in the majority of musculoskeletal conditions.

---

## References (MS-1)

---

**v1.0.2024**

1. Reinus WR. Clinician's guide to diagnostic imaging. New York, NY: Springer Science; 2014.
2. Visconti AJ, Biddle J, Solomon M. Follow-up imaging for vertebral osteomyelitis a teachable moment. *JAMA*. 2014;174(2):184. doi: 10.1001/jamainternmed.2013.12742.
3. Fabiano V, Franchino G, Napolitano M, Ravelli A, Dilillo D, Zuccotti GV. Utility of magnetic resonance imaging in the follow-up of children affected by acute osteomyelitis. *Curr Pediatr Res*. 2017;21(2):354-358.



# Imaging Techniques (MS-2)

---

## Guideline

---

Plain X-Ray (MS-2.1)  
MRI or CT (MS-2.2)  
Ultrasound (MS-2.3)  
Contrast Issues (MS-2.4)  
Positron Emission Tomography (PET/CT) (MS-2.5)  
Nuclear Medicine (MS-2.6)  
References (MS-2)

## Plain X-Ray (MS-2.1)

**MS.IM.0002.1.A**

**v1.0.2024**

- The results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider of the advanced imaging study for all musculoskeletal conditions, unless otherwise noted in the guidelines, to rule out those situations that do not often require advanced imaging, such as: osteoarthritis, acute/healing fracture, dislocation, osteomyelitis, acquired/congenital deformities, and tumors of bone amenable to biopsy or radiation therapy (in known metastatic disease), etc.

## MRI or CT (MS-2.2)

MS.IM.0002.2.C

v1.0.2024

- Magnetic Resonance Imaging (MRI) is often the preferred advanced imaging modality in musculoskeletal conditions because it is superior in imaging the soft tissues and can also define physiological processes in some instances [e.g. edema, loss of circulation (AVN), and increased vascularity (tumors)].
- Computed Tomography (CT) is preferred for imaging cortical bone anatomy; thus, it is useful for studying complex fractures (particularly of the joints), dislocations, and assessing delayed union or non-union of fractures, if plain X-rays are equivocal. CT may be the procedure of choice in individuals who cannot undergo an MRI, such as those with pacemakers.

### **Positional MRI**

- Positional MRI is also referred to as dynamic, standing, weight-bearing, or kinetic MRI. Currently, there is inadequate scientific evidence to support the medical necessity of this study. As such, it should be considered experimental or investigational.

### **Positional CT**

- Positional CT, also referred to as weight-bearing or cone beam CT, may be useful in imaging of the foot and ankle.
  - If a request for foot or ankle imaging with positional CT meets medical necessity criteria for standard CT imaging (as defined in the condition-specific guidelines), the request may be approved.
    - Positional CT of anatomic areas other than the foot and ankle are considered experimental or investigational.

### **dGEMRIC Evaluation of Cartilage**

- Delayed gadolinium enhanced Magnetic Resonance Imaging of Cartilage (dGEMRIC) is a technique where an MRI estimates joint cartilage glycosaminoglycan content after penetration of the contrast agent in order to detect cartilage breakdown. Currently, there is inadequate scientific evidence to support the medical necessity of this study. As such, it should be considered experimental or investigational for the diagnosis and surveillance of, or preoperative planning related to chondral pathology.

---

## Ultrasound (MS-2.3)

---

**MS.IM.0002.3.A**

**v1.0.2024**

- Ultrasound (US) uses sound waves to produce images that can be used to evaluate a variety of musculoskeletal disorders. As with US in general, musculoskeletal US is highly operator-dependent, and proper training and experience are required to perform consistent, high quality evaluations.

## Contrast Issues (MS-2.4)

MS.IM.0002.4.A

v1.0.2024

- Most musculoskeletal imaging (MRI or CT) is without contrast; however, the following examples may be considered with contrast:
  - Tumors, osteomyelitis, and soft tissue infection (without and with contrast)
  - MRI arthrography (with contrast only)
  - MRI for rheumatoid arthritis and inflammatory arthritis (contrast as requested)
  - For individuals with a contrast contraindication, if the advanced imaging recommendation specifically includes contrast, the corresponding advanced imaging study without contrast may be approved as an alternative, although the non-contrast study may not provide an adequate evaluation of the condition of concern.

# Positron Emission Tomography (PET/CT) (MS-2.5)

---

MS.IM.0002.5.A

v1.0.2024

- PET/CT is a nuclear medicine/computed tomography (CT) fusion study that uses a positron emitting radiotracer to create cross-sectional and volumetric images based on tissue metabolism. PET imaging fusion with CT allows for better anatomic localization of the areas of abnormal increased tissue activity seen on PET.
- PET/CT is indicated for imaging of certain musculoskeletal conditions when MRI or CT is equivocal or cannot be performed. See: **Nuclear Medicine (MS-28)** for specific indications.
  - At this time, FDG is the only indicated radiotracer for use with PET/CT in the imaging of musculoskeletal conditions.

## Nuclear Medicine (MS-2.6)

MS.IM.0002.6.A

v1.0.2024

- A bone scan is a nuclear medicine imaging study in which an amount of radioactive material is injected and images are obtained at different time intervals, depending on the condition. A bone scan is done to reveal problems with bone metabolism. Areas where bone cells are repairing themselves show the most activity. It can help diagnose a number of bone conditions, including cancer of the bone or metastasis, location of bone inflammation, fracture, and bone infection.
- Nuclear Medicine WBC Scan is performed using radioactive material which is tagged to the white blood cells. When injected into the body, the material attaches to sites of inflammation/infection. Once distributed in these areas, the sites of suspected infection/inflammation can be seen on nuclear imaging equipment. These can be imaged as a planar study, SPECT study, or SPECT/CT study.
- Bone Marrow Imaging is used in combination with a WBC Scan to help differentiate between true infection and physiological marrow uptake. The bone marrow scan provides a map of the normal physiological white cell uptake that is then compared to the white blood cell scan. Any discordance in white cell uptake (e.g., more WBC uptake than marrow uptake) between the two studies indicates a focus of infection.
- See: **Nuclear Medicine (MS-28)** and condition-specific guidelines for specific indications.

## References (MS-2)

v1.0.2024

1. DeMuro JP, Simmons S, Smith K, et al. Utility of MRI in blunt trauma patients with a normal cervical spine CT and persistent midline neck pain on palpation. *Global Journal of Surgery*. 2013;1(1):4-7. doi:10.12691/js-1-1-2.
2. Hsu W, Hearty TM. Radionuclide imaging in the diagnosis and management of orthopaedic disease. *J Am Acad Orthop Surg*. 2012;20(3):151-159. doi:10.5435/JAAOS-20-03-151.
3. Kayser R, Mahlfeld K, Heyde CE. Partial rupture of the proximal Achilles tendon: a differential diagnostic problem in ultrasound imaging. *Br J Sports Med*. 2005;9(11):838–842. doi:10.1136/bjsm.2005.018416.
4. Ward RJ, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute hip pain-suspected fracture. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Revised: 2018. <https://acsearch.acr.org/docs/3082587/Narrative/>.
5. Mosher TJ, Kransdorf MJ, Adler R, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute trauma to the ankle. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Revised: 2020. <https://acsearch.acr.org/docs/69436/Narrative/>.
6. Small KM, Adler RS, Shah SH, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Atraumatic. *Am Coll Radiol (ACR)*; New 2018. <https://acsearch.acr.org/docs/3101482/Narrative/>.
7. Amini B, Beckmann NM, Beaman FD, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Traumatic. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69433/Narrative/>.
8. Hayes CW, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic elbow pain. *Am Coll Radiol (ACR)*; Date of Origin:1998. Revised: 2022. <https://acsearch.acr.org/docs/69423/Narrative/>.
9. Wise JN, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic foot pain. *Am Coll Radiol (ACR)*; Date of Origin:1998. Revised: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
10. Mintz DN, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic hip pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
11. Rubin DA, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic wrist pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
12. Bennett DL, Nelson JW, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® nontraumatic knee pain. *Am Coll Radiol (ACR)*;1995. Revised: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
13. Murphey MD, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® osteonecrosis of the hip. *Am Coll Radiol (ACR)*;Date of Origin: 1995. Revised: 2022. <https://acsearch.acr.org/docs/69420/Narrative/>.
14. Bruno MA, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® acute hand and wrist trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
15. Bencardino JT, Stone TJ, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® stress (fatigue/insufficiency) fracture, including sacrum, excluding other vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
16. Luchs JS, Flug JA, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic ankle pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.



17. Beaman FD, von Herrmann PF, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> suspected osteomyelitis, septic arthritis, or soft tissue infection (excluding spine and diabetic foot. *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/%203094201/Narrative/>.
18. Kransdorf MJ, Weissman BN, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> suspected osteomyelitis of the foot in patients with diabetes mellitus. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2019. <https://acsearch.acr.org/docs/69340/Narrative/>.
19. Zoga AC, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> soft-tissue masses. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
20. Morrison WB, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> primary bone tumors. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2019. <https://acsearch.acr.org/docs/69421/Narrative/>.
21. Weissman BN, Palestro CJ, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> imaging after total hip arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin:1998. Revised: 2015. <https://acsearch.acr.org/docs/3094200/Narrative/>.
22. Hochman MG, Melenevsky YV, Metter DF, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> imaging after total knee arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69430/Narrative/>.
23. Gyftopoulos S, Rosenberg ZS, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> imaging after shoulder arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2021<https://acsearch.acr.org/docs/3097049/Narrative/>.
24. Patel ND, Broderick DF, Burns J, et al. Expert Panel on Neurologic Imaging. ACR Appropriateness Criteria<sup>®</sup>: low back pain. *Am Coll Radiol (ACR)*; Date of Origin:1996. Last Review: 2021. <https://acsearch.acr.org/docs/69483/Narrative/>.
25. Shetty VS, Reis MN, Aulino JM, et al. Expert Panel on Neurologic Imaging. ACR Appropriateness Criteria<sup>®</sup>: head trauma. *Am Coll Radiol (ACR)*; Date of Origin:1996. Last Review: 2020. <https://acsearch.acr.org/docs/69481/Narrative/>.
26. Li X, Yi P, Curry EJ, Murakami AM. Ultrasonography as a diagnostic, therapeutic, and research tool in orthopaedic surgery. *J Am Acad Orthop Surg*. 2018;26(6):187-196. doi: 10.5435/JAAOS-D-16-00221.
27. de Cesar Netto C, Myerson MS, Day J, et. al. Consensus for the use of weightbearing CT in the assessment of progressive collapsing foot deformity. *Foot Ankle Int*. 2020;41(10):1277-1282.
28. Conti MS, Ellis SJ. Weight-bearing ct scans in foot and ankle surgery. *J Am Acad Orthop Surg*. 2020;28(14):e595-e603.
29. de Cesar Netto C, Schon LC, Thawait GK, et. al. Flexible adult acquired flatfoot deformity: comparison between weight-bearing and non-weight-bearing measurements using cone-beam computed tomography. *J Bone Joint Surg Am*. 2017;99(18):e98.

# 3D Rendering (MS-3)

---

## Guideline

---

3D Rendering (MS-3)  
Reference (MS-3)

## 3D Rendering (MS-3)

MS.TD.0003.U

v1.0.2024

- Indications for musculoskeletal 3-D image post-processing for preoperative planning when conventional imaging is insufficient for:
  - Complex fractures (comminuted or displaced)/dislocations of any joint.
  - Spine fractures, pelvic/acetabulum fractures, intra-articular fractures.
  - Preoperative planning for other complex surgical cases.
- The code assignment for 3-D rendering depends upon whether the 3-D post-processing is performed on the scanner workstation (CPT<sup>®</sup> 76376) or on an independent workstation (CPT<sup>®</sup> 76377).
  - 2-D reconstruction (i.e. reformatting axial images into the coronal plane) is considered part of the tomography procedure, is not separately reportable, and does not meet the definition of 3-D rendering.
  - It is not appropriate to report 3-D rendering in conjunction with CTA and MRA because those procedure codes already include the post-processing.
  - In addition to the term "3-D," the following terms may also be used to describe 3-D post-processing:
    - Maximum intensity projection (MIP)
    - Shaded surface rendering
    - Volume rendering
  - Additionally - If multiple CPT codes are performed for the same indication on the same day, one 3D rendering code is required. If they are performed on separate days, 3D rendering codes are required for each study on each day.
- The 3-D rendering codes require concurrent supervision of image post-processing 3-D manipulation of volumetric data set and image rendering.

## Reference (MS-3)

**v1.0.2024**

1. Bruno MA, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® acute hand and wrist trauma. Am Coll Radiol (ACR); Date of Origin:1995. Revised: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.

# Avascular Necrosis (AVN)/ Osteonecrosis (MS-4)

---

## Guideline

---

AVN (MS-4.1)

References (MS-4)

# AVN (MS-4.1)

MS.AN.0004.1.A

v1.0.2024

- MRI without contrast, MRI without and with contrast, or CT without contrast of the area of interest can be performed when plain x-ray findings are negative or equivocal and clinical symptoms warrant further investigation for suspected avascular necrosis.
- Advanced imaging for AVN confirmed by plain x-ray is appropriate for treatment planning in the following situations:
  - Femoral head:
    - MRI Hip without contrast (CPT<sup>®</sup> 73721) or CT Hip without contrast (CPT<sup>®</sup> 73700)
  - Distal Femur:
    - MRI Knee without contrast (CPT<sup>®</sup> 73721) or CT Knee without contrast (CPT<sup>®</sup> 73700)
  - Talus:
    - MRI Ankle without contrast (CPT<sup>®</sup> 73721) or CT Ankle without contrast (CPT<sup>®</sup> 73700)
  - Tarsal navicular (Kohler Disease):
    - MRI Foot without contrast (CPT<sup>®</sup> 73718) or CT Foot without contrast (CPT<sup>®</sup> 73700)
  - Metatarsal head (Frieberg's Infraction):
    - MRI Foot without contrast (CPT<sup>®</sup> 73718) or CT Foot without contrast (CPT<sup>®</sup> 73700)
  - Humeral head:
    - MRI Shoulder without contrast (CPT<sup>®</sup> 73221) or CT Shoulder without contrast (CPT<sup>®</sup> 73200)
  - Lunate (Kienbock's Disease)/Scaphoid (Preiser's Disease):
    - CT Wrist without contrast (CPT<sup>®</sup> 73200) or MRI Wrist without contrast (CPT<sup>®</sup> 73221)
- Individuals with acute lymphoblastic leukemia and known or suspected osteonecrosis should be imaged according to guidelines in **Acute Lymphoblastic Leukemia (PEDONC-3.2)**
- Known or suspected osteonecrosis in long-term cancer survivors should be imaged according to guidelines in **Osteonecrosis in Long Term Cancer Survivors (PEDONC-19.4)**

## ***Background and Supporting Information***

- Classification systems use a combination of plain radiographs, MRI, and clinical features to stage avascular necrosis.

## References (MS-4)

**v1.0.2024**

1. Calder JD, Hine AL, Pearse MF, Revell PA. The relationship between osteonecrosis of the proximal femur identified by MRI and lesions proven by histological examination. *J Bone Joint Surg Br.* 2008;90(2):154-158.
2. Karantanas AH, Drakonaki EE. The role of MR imaging in avascular necrosis of the femoral head. *Semin Musculoskelet Radiol.* 2011;15(3):281-300. doi:10.1055/s-0031-1278427.
3. Karim AR, Cherian JJ, Jauregui JJ, et al. Osteonecrosis of the knee: review. *Ann Transl Med.* 2015;3(1). doi:10.3978/j.issn.2305-5839.2014.11.13.
4. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic hip pain. *Am Coll Radiol (ACR)*; Revised:2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
5. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic wrist pain. *Am Coll Radiol (ACR)*; Revised:2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
6. Bennett DL, Nelson JW, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® nontraumatic knee pain. *Am Coll Radiol (ACR)*; Date of Origin:1995. Revised: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
7. Murphey MD, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Osteonecrosis. *Am Coll Radiol (ACR)*; Date of Origin:1995. Revised: 2022. <https://acsearch.acr.org/docs/69420/Narrative/>.

# Fractures (MS-5)

---

## Guideline

---

Acute Fracture (MS-5.1)

Suspected Occult/Stress/Insufficiency Fracture/Stress Reaction and Shin Splints (MS-5.2)

Other Indications (MS-5.3)

References (MS-5)



## Acute Fracture (MS-5.1)

MS.FX.0005.1.A

v1.0.2024

- CT or MRI without contrast if **ANY** of the following:
  - Complex (comminuted or displaced) fracture with or without dislocation on plain x-ray.
    - CT is preferred unless it is associated with neoplastic disease when MRI without/with contrast is preferred unless MRI contraindicated.
  - Individual presents initially to the requesting provider with a documented history of an acute traumatic event at least two weeks prior with a negative plain x-ray at the time of this face-to-face encounter and a clinical suspicion for an occult/stress/insufficiency fracture see: **Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints (MS-5.2)**.
- For osteochondral fracture or osteochondral injury, see: **Chondral/Osteochondral Lesions, Including Osteochondritis Dissecans and Fractures (MS-13.1)**

# Suspected Occult/Stress/Insufficiency Fracture/Stress Reaction and Shin Splints (MS-5.2)

MS.FX.0005.2.A

v1.0.2024

- MRI without contrast can be performed for suspected hip/femoral neck, tibia, pelvis/sacrum, tarsal navicular, proximal fifth metatarsal, or scaphoid occult/stress/insufficiency fractures, and suspected atypical femoral shaft fractures related to bisphosphonate use if the initial evaluation of history, physical exam and plain x-ray fails to establish a definitive diagnosis.
  - CT without contrast can be performed as an alternative to MRI for suspected occult/insufficiency fractures of the pelvis/hip and suspected atypical femoral shaft fractures related to bisphosphonate see: **Pelvis (MS-23)** and **Hip (MS-24)**, and suspected occult fractures of the scaphoid see: **Wrist (MS-21)**.
  - For suspected fractures, when MRI cannot be performed, any one of the following is indicated:
    - Tc-99m Bone scan whole body (CPT<sup>®</sup> 78306) with SPECT of the area of interest (CPT<sup>®</sup> 78803) OR
    - Bone scan (CPT<sup>®</sup> 78315, 78300) OR
    - SPECT/CT (CPT<sup>®</sup> 78830)
    - See: **Nuclear Medicine (MS-28)**
- MRI or CT without contrast can be performed for all other suspected occult/stress/insufficiency fractures with either of the following:
  - Repeat plain x-rays remain non-diagnostic for fracture after a minimum of 10 days of provider-directed conservative treatment, or
  - Initial plain x-rays obtained a minimum of 14 days after the onset of symptoms are non-diagnostic for fracture
  - For suspected fractures, when MRI cannot be performed, any one of the following is indicated:
    - Tc-99m Bone scan whole body (CPT<sup>®</sup> 78306) with SPECT of the area of interest (CPT<sup>®</sup> 78803) OR
    - Bone scan (CPT<sup>®</sup> 78315, 78300) OR
    - SPECT/CT (CPT<sup>®</sup> 78830)
    - See: **Nuclear Medicine (MS-28)**
- MRI of the lower leg without contrast (CPT<sup>®</sup> 73718) for suspected shin splints when **BOTH** of the following are met:
  - Initial plain x-ray

- Failure of a 6-week trial of provider-directed conservative treatment
- For stress reaction, advanced imaging is not medically necessary for surveillance or “return to play” decisions regarding a stress reaction identified on an initial imaging study.
- MRI without contrast of the area of interest for stress fracture follow-up imaging for “return to play” evaluation at least 3 months after the initial imaging study for stress fracture.
- For periprosthetic fractures related to joint replacement see: **Post-Operative Joint Replacement Surgery (MS-16.1)**, **Shoulder (MS-19)**, **Elbow (MS-20)**, **Hip (MS-24)**, **Knee (MS-25)**, and **Ankle (MS-26)**.

## Other Indications (MS-5.3)

MS.FX.0005.3.A

v1.0.2024

- CT or MRI without contrast after recent (within 30 days) plain x-ray if **ONE** of the following is present:
  - Concern for delayed union or non-union of fracture, osteotomy, or joint fusions.
  - Part of preoperative evaluation for a planned surgery of a complex fracture with or without dislocation.

## References (MS-5)

v1.0.2024

1. Bencardino JT, Stone TJ, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
2. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
3. Bruno MA, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hand and Wrist Trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
4. Luchs JS, Flug JA, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Ankle Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
5. Ward RJ, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hip Pain-Suspected Fracture. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Revised: 2018. <https://acsearch.acr.org/docs/3082587/Narrative/>.
6. Mosher TJ, Kransdorf MJ, Adler R, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Ankle. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Revised: 2020. <https://acsearch.acr.org/docs/69436/Narrative/>.
7. Hayes CW, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Elbow Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2022. <https://acsearch.acr.org/docs/69423/Narrative/>.
8. Wise JN, Weissman BN, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
9. Greene WB. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:568-570.
10. Galbraith RM, Lavalley ME. Medial tibial stress syndrome: conservative treatment options. *Curr Rev Muscuolskelet Med*. 2009;2:127-133. doi:10.1007/s12178-009-9055-6.
11. Boks SS, Vroegindewij D, Kroes BW, et al. MRI follow-up of posttraumatic bone bruises of the knee in general practice. *AJR Am J Roentgenol*. 2007;189 556-562. doi:10.2214/AJR.07.2276.
12. Kaeding CC, Yu JR, Wright R, et al. Management and return to play of stress fractures. *Clin J Sport Med*. 2005;15:442-7.
13. Sormaala MJ, Niva MH, Kiuru MJ, et al. Stress injuries of the calcaneus detected with magnetic resonance imaging in military recruits. *J Bone Joint Surg Am*. 2006;88:2237-42. doi:10.2106/JBJS.E.01447.
14. Shin AY, Morin WD, Gorman JD, et al. The superiority of magnetic resonance imaging in differentiating the cause of hip pain in endurance athletes. *Am J Sports Med*. 1996;24:168-76. doi:10.1177/036354659602400209.
15. Slocum KA, Gorman JD, Puckett ML, et al. Resolution of abnormal MR signal intensity in patients with stress fractures of the femoral neck. *AJR Am J Roentgenol*. 1997;168:1295-9. doi:10.2214/ajr.168.5.9129429.
16. Fredericson M, Bergman AG, Hoffman KL, et al. Tibial stress reaction in runners. Correlation of clinical symptoms and scintigraphy with a new magnetic resonance imaging grading system. *Am J Sports Med*. 1995;23(4):472-81. doi:10.1177/036354659502300418.
17. Bernstein EM, Kelsey TJ, Cochran GK, Deafenbaugh BK, Kuhn KM. Femoral neck stress fractures: An updated review. *J Am Acad Orthop Surg*. 2022;30:302-311. doi:10.5435/JAAOS-D-21-00398.

# Foreign Body (MS-6)

---

## Guideline

---

Foreign Body – General (MS-6.1)

References (MS-6)

## Foreign Body – General (MS-6.1)

**MS.FB.0006.1.A**

**v1.0.2024**

- Ultrasound (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) or CT without contrast or MRI without and with contrast or MRI without contrast of the area of interest can be approved after plain x-rays rule out the presence of radiopaque foreign bodies.
  - Ultrasound (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) is the preferred imaging modality for radiolucent (non-radiopaque) foreign bodies (e.g. wood, plastic).
  - CT without contrast is recommended when plain x-rays are negative and a radiopaque foreign body is still suspected, as CT is favored over MRI for the identification of foreign bodies
  - MRI without and with contrast is an alternative to US and CT for assessing the extent of infection associated with a suspected foreign body

## References (MS-6)

**v1.0.2024**

1. Bancroft LW, Kransdorf MJ, Adler R, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Acute Trauma to the Foot. *Am Coll Radiol (ACR)*; Date of Origin: 2010. Revised: 2019. <https://acsearch.acr.org/docs/70546/Narrative/>.
2. Beaman FD, von Herrmann PF, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/%203094201/Narrative>.
3. Chan C, Salam GA. Splinter removal. *Am Fam Physician*. 2003;67(12):2557-2562.
4. Peterson JJ, Bancroft LW, Kransdorf MJ. Wooden foreign bodies: imaging appearance. (*AJR Am J Roentgenol*. 2002;178(3):557-562. doi:10.2214/ajr.178.3.1780557.
5. Jarraya M, Hayashi D, de Villiers RV, et al. Multimodality imaging of foreign bodies of the musculoskeletal system. (*AJR Am J Roentgenol*. 2014;203(1):W92-102. doi:10.2214/AJR.13.11743.



# Ganglion Cysts (MS-7)

---

## Guideline

---

Ganglion Cysts – General (MS-7.1)

References (MS-7)

## **Ganglion Cysts – General (MS-7.1)**

**MS.GC.0007.1.A**

**v1.0.2024**

- Plain x-ray is the initial imaging study for ganglion cysts.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- MRI without contrast or MRI without and with contrast or US (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) is appropriate for surgical planning.
- Advanced imaging is not indicated for ganglions that can be diagnosed by history and physical examination.

## References (MS-7)

**v1.0.2024**

1. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging ACR Appropriateness Criteria<sup>®</sup> Chronic wrist pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
2. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Soft-Tissue Masses. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
3. Freire V, Guerini H, Campagna R, et al. Imaging of hand and wrist cysts: a clinical approach. (*AJR Am J R Roentgenol*. 2012;199(5):W618-W628. doi:10.2214/AJR.11.8087.
4. Vo P, Wright T, Hayden F, Dell P, et al. Evaluating dorsal wrist pain: MRI diagnosis of occult dorsal wrist ganglion. *J Hand Surg Am*. 1995;20(4):667-670. doi:10.1016/S0363-5023(05)80288-6.
5. Teefey SA, Dahiya N, Middleton WD, et al. Ganglia of the hand and wrist: a sonographic analysis. *AJR Am J Roentgenol*. 2008;191(3):716-720. doi:10.2214/AJR.07.3438.

# Gout/Calcium Pyrophosphate Deposition Disease (CPPD)/ Pseudogout/ Chondrocalcinosis (MS-8)

---

## Guideline

---

Gout – General (MS-8.1)

CPPD (Pseudogout/Chondrocalcinosis) – General (MS-8.2)

References (MS-8)

## Gout – General (MS-8.1)

MS.GD.0008.1.A

v1.0.2024

- CT without contrast, MRI without contrast, or MRI without and with contrast of the area of interest is indicated when **BOTH** of the following are met:
  - Initial plain x-ray to rule out other potential disease processes
  - Infection or neoplasm is in the differential diagnosis for soft-tissue tophi

### ***Background and Supporting Information***

- Early stages of gout can be diagnosed clinically since radiographic findings are not present early in the disease course.

# CPPD (Pseudogout/Chondrocalcinosis) – General (MS-8.2)

---

MS.GD.0008.2.A

v1.0.2024

- Calcium pyrophosphate deposition disease (CPPD), also called pseudogout, can often be diagnosed from plain x-rays; advanced diagnostic imaging is generally not medically necessary.

## References (MS-8)

v1.0.2024

1. Hsu CY, Shih TT, Huang KM, et al. Tophaceous gout of the spine: MR imaging features. *Clin Radiol*. 2002;57(10):919.
2. Schumacher HR Jr, Becker MA, Edwards NL, et al. Magnetic resonance imaging in the quantitative assessment of gouty tophi. *Int J Clin Pract*. 2006;60(4):408. doi:10.1111/j.1368-5031.2006.00853.x.
3. McQueen FM, Doyle A, Reeves Q, Gao A. Bone erosions in patients with chronic gouty arthropathy are associated with tophi but not bone oedema or synovitis: new insights from a 3 T MRI study. *Rheumatology*. 2014;53(1):95-103. doi:10.1093/rheumatology/ket329.
4. Dore RK. Gout: What primary care physicians want to know. *J Clin Rheumatol*. 2008;14(5S):S47-S54. doi:10.1097/RHU.0b013e3181896c35.
5. Eggebeen AT. Gout: an update. *Am Fam Physician*. 2007;76(6):801-808.
6. Burns C, Wortmann RL. Gout. In: Imboden JB, Hellmann DB, Stone JH, eds. *CURRENT Diagnosis & Treatment: Rheumatology*. 3<sup>rd</sup> edition. New York: McGraw-Hill; 2013:332-338.
7. Jacobson JA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Chronic extremity joint pain-suspected inflammatory arthritis. *Am Coll Radiol (ACR)*; 2017;14(5):S81-S89. [http://www.jacr.org/article/S1546-1440\(17\)30183-7/fulltext](http://www.jacr.org/article/S1546-1440(17)30183-7/fulltext).

# Infection/ Osteomyelitis (MS-9)

---

## Guideline

---

Infection – General (MS-9.1)  
Septic Joint (MS-9.2)  
References (MS-9)



## Infection – General (MS-9.1)

MS.OI.0009.1.A

v1.0.2024

- MRI without contrast, MRI without and with contrast, CT without contrast, CT with contrast, or Ultrasound (CPT<sup>®</sup> 76881 or 76882) of the affected area is appropriate after plain x-ray(s) in the following scenarios:
  - Plain x-ray(s) do not demonstrate infection, **AND**
  - Plain x-ray(s) do not suggest alternative diagnoses such as neuropathic arthropathy or fracture, **AND**
  - Soft tissue or bone infection (osteomyelitis) is suspected **OR**
  - Plain x-ray(s) are positive for infection, **AND**
  - The extent of infection into the soft tissues and any skip lesions require evaluation
- Individuals with suspected spinal infections
  - See: **Red Flag Indications (SP-1.2)** for advanced imaging guidelines
- Individuals with diabetic foot infections after plain x-ray(s)
  - See: **Foot (MS-27)** for advanced imaging guidelines
- For nuclear medicine studies appropriate in specific scenarios, see: **Nuclear Medicine (MS-28)**

## Septic Joint (MS-9.2)

MS.OI.0009.2.A

v1.0.2024

- MRI without and with contrast, MRI without contrast, CT without contrast, CT with contrast, or Ultrasound (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) of the affected joint is appropriate when standard or image-guided arthrocentesis is contraindicated, unsuccessful, or non-diagnostic, and the clinical documentation satisfies ALL of the following criteria:
  - History and physical examination findings [One of the following]:
    - Development of an acutely hot and swollen joint (< 2 weeks)
    - Decreased range of motion due to pain
    - Documented fever
  - Laboratory tests [One of the following]:
    - Leukocytosis
    - Elevated ESR or C-reactive protein
    - Analysis of the joint fluid is non-diagnostic
  - Plain x-ray of the joint
    - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- MRI without and with contrast, MRI without contrast, CT without contrast, or CT with contrast of the affected joint is appropriate after plain x-rays if the arthrocentesis is diagnostic and if there is a confirmed septic joint, to evaluate the extent of infection into the soft tissues and any skip lesions that would require evaluation.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider

### ***Background and Supporting Information***

- Analysis of joint fluid is most often sufficient to diagnose a septic joint.

## References (MS-9)

v1.0.2024

1. Coakley G, Mathews C, Field M, et al. BSR & BHP, BOA, RCGP and BSAC guidelines for management of the hot swollen joints in adults. *Rheumatology*. 2006;45(8):1039-1041. doi:10.1093/rheumatology/ke1163a.
2. Karchevsky M, Schweitzer ME, Morrison WB, et al. MRI findings of septic arthritis and associated osteomyelitis in adults. (*AJR*) *Am J Roentgenol*. 2004;182(1):119-122. doi:10.2214/ajr.182.1.1820119.
3. Griffin LY. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:918.
4. Staheli LT. Septic arthritis. In: Staheli LT, ed. *Fundamentals of Pediatric Orthopedics*. 4<sup>th</sup> edition. Philadelphia, PA: Lippincott Williams & Wilkins; 2008:110-111.
5. Kransdorf MJ, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Suspected osteomyelitis of the foot in patients with diabetes mellitus. *Am Coll Radiol (ACR)*. 2012. <https://acsearch.acr.org/docs/69340/Narrative/>.
6. Beaman FD, von Herrmann PF, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Suspected osteomyelitis, septic arthritis, or soft tissue infection (excluding spine and diabetic foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/3094201/Narrative/>.
7. Rubin DA, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Chronic wrist pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
8. Small KM, Adler RS, Shah SH, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Shoulder Pain - Atraumatic. *Am Coll Radiol (ACR)*; New 2018. <https://acsearch.acr.org/docs/3101482/Narrative/>.
9. Amini B, Beckmann NM, Beaman FD, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Shoulder Pain - Traumatic. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69433/Narrative/>.
10. Mintz DN, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Chronic hip pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
11. Reinus WR. *Clinician's Guide to Diagnostic Imaging*. 2014. Springer-Verlag New York.
12. Visconti AJ, Biddle J, Solomon M. Follow-up imaging for vertebral osteomyelitis a teachable moment. *JAMA Intern Med*. 2014;174(2):184. doi:10.1001/jamainternmed.2013.12742.
13. Fabiano V, Franchino G, Napolitano M, et. al. Utility of magnetic resonance imaging in the follow-up of children affected by acute osteomyelitis. *Curr Pediatr Res*. 2017;21(2):354-358.
14. Patel ND, Broderick DF, Burns J, et. al. Expert Panel on Neurologic Imaging. ACR Appropriateness Criteria<sup>®</sup>: Low back pain. *Am Coll Radiol (ACR)*. 2015; Revised: 2021. <https://acsearch.acr.org/docs/69483/Narrative/>.

# Soft Tissue Mass or Lesion of Bone (MS-10)

---

## Guideline

---

Soft Tissue Mass (MS-10.1)

Lesion of Bone (MS-10.2)

References (MS-10)

## Soft Tissue Mass (MS-10.1)

### MS.ST.0010.1.A

v1.0.2024

- History and physical exam should include documentation of: location, size, duration, growing or stable, solid/cystic, fixed/not fixed to the bone, discrete or ill-defined, and an association with pain.
- Plain x-ray is indicated as the initial imaging study, with the exception of individuals with cancer predisposition syndrome.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- MRI without and with contrast or without contrast or US of the area of interest (CPT<sup>®</sup> 76881 or 76882) is appropriate when ANY of the following are met after plain x-ray:
  - Soft tissue mass(es)
  - Surgical planning
- Known or suspected soft tissue mass in an individual with a cancer predisposition syndrome, see **Screening Imaging in Cancer Predisposition Syndromes (PEDONC-2)** in the Pediatric and Special Populations Oncology Imaging Guidelines.
- CT with contrast or CT without and with contrast is appropriate when MRI is contraindicated or after a metal limiting MRI evaluation.
- Advanced imaging is not indicated for:
  - Subcutaneous lipoma with no surgery planned
  - Ganglia, see: **Ganglion Cysts (MS-7)**
  - Sebaceous cyst

### **Background and Supporting Information**

- Plain x-rays can determine if an advanced imaging procedure is indicated, and if so, which modality is most appropriate. If non-diagnostic, these initial plain x-rays can provide complementary information if advanced imaging is indicated.

## Lesion of Bone (MS-10.2)

MS.ST.0010.2.A

v1.0.2024

- History and physical exam should include documentation of: location, size, duration, growing or stable, discrete or poorly defined, and an association with pain.
- Complete radiograph of the entire bone containing the lesion of bone is required prior to consideration of advanced imaging. Many benign bone tumors have a characteristic appearance on plain x-ray and advanced imaging is not necessary.
- MRI without and with contrast, MRI without contrast, or CT without contrast may be indicated if ONE of the following applies:
  - Diagnosis uncertain based on plain x-ray appearance
  - Imaging requested for preoperative planning
- MRI without and with contrast or without contrast is appropriate when plain x-ray reveals an osteochondroma with clinical concern of malignant transformation.
- For Paget's Disease:
  - Bone scan (See: **Nuclear Medicine (MS-28)**) **OR**
  - MRI (contrast as requested) can be considered if the diagnosis (based on plain x-rays and laboratory studies) is in doubt.
  - MRI (contrast as requested) can be considered if malignant degeneration, which occurs in up to 10% of cases, is suspected.

## References (MS-10)

**v1.0.2024**

1. Peterson JJ, Beaman FD, Fox MG, et al. ACR Practice Guideline. ACR-SSR Practice Guideline for the Performance and Interpretation of Magnetic Resonance Imaging (MRI) of bone and soft tissue tumors. *Am Coll Radiol*. Revised 2020. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-SoftTissue-Tumors.pdf>.
2. Zoga AC, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Soft-tissue masses. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
3. Hayes CW, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic elbow pain. *Am Coll Radiol (ACR)*; Revised 2022. <https://acsearch.acr.org/docs/69423/Narrative/>.
4. Musculoskeletal Tumor Society: Systematic Literature Review on the Use of Imaging Prior to Referral to a Musculoskeletal Oncologist. Rosemont, IL, Musculoskeletal Tumor Society, February 2018.
5. Schneider D, Hofmann MR, and Peterson JA. Diagnosis and treatment of Paget's Disease of Bone. *Am Fam Physician*. 2002;65(10):2069-2072. <https://www.aafp.org/afp/2002/0515/p2069.html>.
6. Theodorou DJ, Theodorou SJ, and Kakitsubata Y. Imaging of Paget Disease of bone and its musculoskeletal complications: review. (*AJR*) *Am J Roentgenol*. 2012;196(6):S64-S75.
7. Sinha S and Peach AH. Diagnosis and management of soft tissue sarcoma. *BMJ*. 2010;341:c7170. doi:10.1136/bmj.c7170.
8. Wu JS, Hochman MG. Soft-tissue tumors and tumorlike lesions: a systematic imaging approach. *Radiology*. 2009;253(2):297-316. doi:10.1148/radiol.2532081199.

# Muscle/Tendon Unit Injuries/Diseases (MS-11)

---

## Guideline

---

Muscle/Tendon Unit Injuries/Diseases (MS-11.1)  
Acute Compartment Syndrome (MS-11.2)  
Chronic Exertional Compartment Syndrome (MS-11.3)  
References (MS-11)



# Muscle/Tendon Unit Injuries/Diseases (MS-11.1)

---

MS.MI.0011.1.A

v1.0.2024

- Plain x-ray is the initial imaging study for muscle/tendon unit injuries.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- MRI without contrast or US (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) is supported for **EITHER** of the following:
  - Suspected partial tendon rupture of a specific (named) tendon
  - Complete tendon rupture of a specific named tendon for preoperative planning
- MRI is not medically necessary for muscle belly strains/muscle tears
- See: **Shoulder (MS-19)** for clinical suspicion of a partial or complete rotator cuff tear
- See: **Inflammatory Muscle Diseases (PN-6.2)** in the Peripheral Nerve Disorders Imaging Guidelines and **Inflammatory Muscle Diseases (PEDMS-10.3)** in the Pediatric Musculoskeletal Imaging Guidelines

# **Acute Compartment Syndrome (MS-11.2)**

**MS.MI.0011.2.A**

**v1.0.2024**

- Advanced imaging is not indicated. Diagnosis is made clinically and by direct measurement of compartment pressure and is a surgical emergency.

## ***Background and Supporting Information***

- Noninvasive methods of measuring compartment pressures and diagnosing acute compartment syndrome are under study, but are currently experimental, investigational, and unproven.

# Chronic Exertional Compartment Syndrome (MS-11.3)

---

MS.MI.0011.3.A

v1.0.2024

- Advanced imaging should only be considered when ruling out other potential causes of extremity pain following a plain x-ray and conservative treatment as indicated.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider

## ***Background and Supporting Information***

- Direct measurement of compartment pressure remains the diagnostic standard. Noninvasive methods of measuring compartment pressures and diagnosing chronic exertional compartment syndrome are under study, but are currently experimental, investigational, and unproven.

## References (MS-11)

**v1.0.2024**

1. Luchs JS, Flug JA, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic ankle pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
2. Griffin LY. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:452.
3. Kayser R, Mahlfeld K, Heyde CE. Partial rupture of the proximal Achilles tendon: a differential diagnostic problem in ultrasound imaging. *Br J Sports Med*. 2005;39:838-842. doi:10.1136/bjism.2005.018416.
4. Rominger MB, Lukosch CJ, Bachmann GF. MR imaging of compartment syndrome of the lower leg: a case control study. *Eur Radiol*. 2004;14:1432-1439. doi:10.1007/s00330-004-2305-5.
5. McDonald S, Bearcroft P. Compartment syndromes. *Semin Musculoskelet Radiol*. 2010;14(2):236-244. doi:10.1055/s-0030-1253164.
6. Ringler MD, Litwiller DV, Felmler JP, et al. MRI accurately detects chronic exertional compartment syndrome: a validation study. *Skeletal Radiology*. 2013;42:385-392. doi:10.1007/s00256-012-1487-1.
7. van den Brand JG, Nelson T, Verleisdonk EJ, van der Werken C. The diagnostic value of intracompartmental pressure measurement, magnetic resonance imaging, and near-infrared spectroscopy in chronic exertional compartment syndrome: a prospective study in 50 patients. *Am J Sports Med*. 2005;33:699-704. doi:10.1177/0363546504270565.
8. Heer ST, Callander JW, Kraeutler MJ, Mei-Dan O, Mulcahey MK. Hamstring Injuries. *The Journal of Bone and Joint Surgery*. 2019;101(9):843-853. doi:10.2106/jbjs.18.00261.

# Osteoarthritis (MS-12)

---

## Guideline

---

Osteoarthritis (MS-12.1)

Treatment Planning (Non-Surgical and Surgical, Other Than Joint Replacement)  
(MS-12.2)

Imaging Prior to Non-Customized-to-Patient Joint Replacement Surgery/Not for  
Intraoperative Navigation (MS-12.3)

Customized-to-Patient Joint Replacement Surgery/Intraoperative Navigation (MS-12.4)

References (MS-12)

# Osteoarthritis (MS-12.1)

MS.OT.0012.1.A

v1.0.2024

- Plain x-ray is the initial imaging study for osteoarthritis.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider

## ***Background and Supporting Information***

- Plain x-rays are performed initially and will reveal characteristic joint space narrowing, osteophyte formation, cyst formation, and subchondral sclerosis.

# Treatment Planning (Non-Surgical and Surgical, Other Than Joint Replacement) (MS-12.2)

MS.OT.0012.2.A

v1.0.2024

- Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider, unless otherwise specified below.
- CT without contrast is appropriate when ALL of the following apply:
  - Requested for treatment planning, AND
  - Congenital or significant atypical post-traumatic arthritic deformities are identified on plain x-ray, AND
  - The aforementioned deformities require further evaluation of their clinical significance, AND
  - The request is related to the shoulder, elbow, wrist, hip, knee, or ankle
- MRI Knee without contrast (CPT<sup>®</sup> 73721) is appropriate in an individual with osteoarthritis for clinical suspicion of a symptomatic degenerative meniscus tear following plain x-rays and conservative treatment. See: **Knee (MS-25)**
- MRI arthrogram or CT arthrogram is appropriate when joint sparing/salvage reconstructive surgery is planned for the following:
  - Suspected concomitant rotator cuff tear of the shoulder - See: **Shoulder (MS-19)**
  - Suspected concomitant labral tear of the shoulder - See: **Shoulder (MS-19)**
  - Suspected concomitant labral tear of the hip - See: **Hip (MS-24)**
  - Suspected concomitant internal derangement of the knee - See: **Knee (MS-25)**

# Imaging Prior to Non-Customized-to-Patient Joint Replacement Surgery/Not for Intraoperative Navigation (MS-12.3)

MS.OT.0012.3.A

v1.0.2024

- The following imaging studies are appropriate per the listed criteria after plain x-ray has been performed
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
    - CT without contrast is appropriate when ALL of the following apply:
      - Requested for treatment planning, **AND**
      - Congenital or significant atypical post-traumatic arthritic deformities are identified on plain x-ray, **AND**
      - The aforementioned deformities require further evaluation of their clinical significance, **AND**
      - The request is related to the shoulder, elbow, wrist, hip, knee, or ankle
    - CT Shoulder without contrast (CPT<sup>®</sup> 73200) and/or MRI Shoulder without contrast (CPT<sup>®</sup> 73221) are appropriate for preoperative planning prior to shoulder replacement
    - For the clinical imaging criteria regarding preoperative joint replacement surgery for each anatomic area, refer to the anatomic area tables:
      - **Shoulder (MS-19)**
      - **Elbow (MS-20)**
      - **Wrist (MS-21)**
      - **Hip (MS-24)**
      - **Knee (MS-25)**
      - **Ankle (MS-26)**



# Customized-to-Patient Joint Replacement Surgery/Intraoperative Navigation (MS-12.4)

MS.OT.0012.4.U

v1.0.2024

- The following imaging studies are appropriate per the listed criteria after plain x-ray has been performed.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- CT without contrast or MRI without contrast of the shoulder, elbow, wrist, hip, knee, or ankle is appropriate\* when the request is for:
  - Treatment planning for customized-to-patient joint replacement surgery, OR
  - Surgical planning using intraoperative navigation for joint replacement surgery (e.g. MAKOplasty)
- AND
- The joint replacement surgery has been approved or does not require prior authorization
- \*The preoperative imaging listed above is considered **not medically necessary** if any of the following are deemed not medically necessary, not a covered benefit, or experimental, investigational, or unproven by the health plan:
  - Joint replacement surgery
  - Customized-to-patient implant
  - Computer assisted surgical navigation (e.g. MAKOplasty)
- See: **Unlisted Procedures/Therapy Treatment Planning (Preface-4.3)** in the Preface Imaging Guidelines

## References (MS-12)

**v1.0.2024**

1. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
2. Bennett DL, Nelson JW, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Nontraumatic Knee Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
3. Manek NJ, Lane NE. Osteoarthritis: Current concepts in diagnosis and management. *Am Fam Physician* 2000 March;61(6):1795-1804. <https://www.aafp.org/afp/2000/0315/p1795.html>.
4. Griffin LY. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:84.
5. Quatman CE, Hettrich CM, Schmitt LC, et. al. The Clinical Utility and Diagnostic Performance of MRI for Identification of Early and Advanced Knee Osteoarthritis: A Systematic Review. *Am J Sports Med*. 2011;39(7):1557–1568. doi:10.1177/0363546511407612.
6. Braun HJ, Gold GE. Diagnosis of osteoarthritis: imaging. *Bone*. 2012;51(2):278–288. doi:10.1016/j.bone.2011.11.019.

# Chondral/Osteochondral Lesions (MS-13)

---

## Guideline

---

Chondral/Osteochondral Lesions, Including Osteochondritis Dissecans and Fractures (MS-13.1)

References (MS-13)

# Chondral/Osteochondral Lesions, Including Osteochondritis Dissecans and Fractures (MS-13.1)

MS.OD.0013.1.A

v1.0.2024

- MRI without contrast, MRI with contrast (arthrogram), or CT with contrast (arthrogram) of the joint or area of interest is indicated when **EITHER** of the following are met:
  - Plain x-rays are negative and an osteochondral fracture is still suspected
  - Plain x-ray and clinical exam suggest an unstable osteochondral injury
- If plain x-rays show a non-displaced osteochondral fragment, follow-up imaging should be with plain x-rays. Advanced imaging is not necessary.
- MRI without contrast or CT without contrast is indicated when healing (including post-operative fixation) cannot be adequately assessed on follow-up plain x-rays.
- See anatomical table sections for recommendations on anatomy-specific osteochondral injuries
  - See: **Ankle (MS-26)** for suspected osteochondral injury of the ankle
  - See: **Elbow (MS-20)** for suspected osteochondral injury of the elbow

## References (MS-13)

**v1.0.2024**

1. Bridges MD, Berland LL, Cernigliaro JG, et al. ACR Practice Guideline. ACR-SSR Practice Guideline for the Performance and Interpretation of Magnetic Resonance Imaging (MRI). *Am Coll Radiol (ACR)*. 2017. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/mr-perf-interpret.pdf?la=en>.
2. Bennett DL, Nelson JW, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Nontraumatic Knee Pain. *Am Coll Radiol (ACR)*; 2012. Last Review: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
3. Rubin DA, Anderson MW, Hastreiter DM, et al. ACR Practice Guideline. ACR-SSR Practice Guideline for the Performance and Interpretation of Magnetic Resonance Imaging (MRI) of the elbow. *Am Coll Radiol (ACR)*. Revised 2021. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/mr-elbow.pdf?la=en>.

# Osteoporosis (MS-14)

---

## Guideline

---

Osteoporosis (MS-14)

References (MS-14)

# Osteoporosis (MS-14)

MS.OP.0014.A

v1.0.2024

- Plain x-ray is not required.
- Quantitative CT (CPT<sup>®</sup> 77078) can be approved for screening when DXA scanner is unavailable or known to be inaccurate for ANY of the following populations:
  - Women age ≥65 years
  - Men age >70 years
  - Women age <65 years who have additional risk factors for osteoporosis based on medical history and other findings:
    - Estrogen deficiency
    - A history of maternal hip fracture that occurred after age 50 years
    - Low body mass (<127 lb. or 57.6 kg)
    - History of amenorrhea (>1 year before age 42 years)
  - Women age <65 years or men age <70 years who have additional risk factors:
    - Current use of cigarettes
    - Loss of height, thoracic kyphosis
  - Individuals of any age with bone mass osteopenia or fragility fractures on imaging studies such as radiographs, CT, or MRI
  - Individuals age 50 years and older who develop a wrist, hip, spine, or proximal humerus fracture with minimal or no trauma, excluding pathologic fractures
  - Individuals of any age who develop 1 or more insufficiency fractures
  - Premenopausal females or males age 20 to 50 years with risk factors:
    - Individuals with medical conditions that could alter bone mineral density
      - Chronic renal failure
      - Rheumatoid arthritis and other inflammatory arthritides
      - Eating disorders, including anorexia nervosa and bulimia
      - Organ transplantation
      - Prolonged immobilization
      - Conditions associated with secondary osteoporosis, such as gastrointestinal malabsorption or malnutrition, sprue, osteomalacia, vitamin D deficiency, endometriosis, acromegaly, chronic alcoholism or established cirrhosis, and multiple myeloma
      - Individuals who have had gastric bypass for obesity
      - Individuals with an endocrine disorder known to adversely affect bone mineral density (e.g., hyperparathyroidism, hyperthyroidism, or Cushing syndrome)
  - Individuals receiving (or expected to receive) glucocorticoid therapy for >3 months

- Hypogonadal men older than 18 years and men with surgically or chemotherapeutically-induced castration
- Individuals beginning or receiving long-term therapy with medications known to adversely affect BMD (e.g., anti-convulsant drugs, androgen deprivation therapy, aromatase inhibitor therapy, or chronic heparin)

**Note:** Repeat screening quantitative computed tomography (QCT) can be approved no sooner than every two years.

- Quantitative CT scan (CPT<sup>®</sup> 77078) can be approved for non-screening/monitoring when DXA scanner is unavailable or known to be inaccurate for ANY of the following circumstances:
  - Follow-up in cases where QCT was the original study
  - Multiple healed vertebral compression fractures
  - Significant scoliosis
  - Advanced arthritis of the spine due to increased cortical sclerosis often with large marginal osteophytes. Obese individual over the weight limit of the dual-energy x-ray absorptiometry (DXA) exam table
  - Severely obese individuals (BMI >35kg/m<sup>2</sup>)
  - Extremes in body height (i.e. very large and very small individuals)
  - Individuals with extensive degenerative disease of the spine
  - A clinical scenario that requires sensitivity to small changes in trabecular bone density (parathyroid hormone and glucocorticoid treatment monitoring).

**Note:** Repeat non-screening/monitoring QCT can be approved no earlier than one year following a change in treatment regimen, and only when the results will directly impact a treatment decision.



---

## References (MS-14)

---

**v1.0.2024**

1. American Association of Clinical Endocrinologists (AACE) Menopause Guidelines Revision Task Force. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract.* 2016;22(Suppl 4):1-42. <https://www.aace.com/files/postmenopausal-guidelines.pdf>.
2. Coleman F, de Buer SJ, LeBoff MS, et al. National Osteoporosis Foundation (NOF). Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int.* 2014;25(10):2359–2381. doi:10.1007/s00198-014-2794-2.
3. U.S. Preventive Services Task Force (USPSTF). Final Recommendation Statement Osteoporosis: Screening. January 2011.
4. Ward RJ, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Osteoporosis and Bone Mineral Density. *Am Coll Radiol (ACR)*; Revised 2022. <https://acsearch.acr.org/docs/69358/Narrative/>.

# Rheumatoid Arthritis (RA) and Inflammatory Arthritis (MS-15)

---

## Guideline

---

Rheumatoid Arthritis (RA) and Inflammatory Arthritis (MS-15.1)  
Pigmented Villonodular Synovitis (PVNS) (MS-15.2)  
References (MS-15)

# Rheumatoid Arthritis (RA) and Inflammatory Arthritis (MS-15.1)

MS.RA.0015.1.A

v1.0.2024

- Plain x-ray, physical exam and appropriate laboratory studies\* are required prior to advanced imaging.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- MRI without contrast OR MRI without and with contrast or US (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) is appropriate for the most symptomatic joint, or of the dominant hand or wrist, in **ALL** of the following situations:
  - When diagnosis is uncertain prior to initiation of drug therapy.
  - To study the effects of treatment with disease modifying anti-rheumatic drug (DMARD) therapy.
  - To identify seronegative RA individuals that might benefit from early DMARD therapy.
  - To determine change in treatment, such as:
    - Switching from standard DMARD therapy to tumor necrosis factor (TNF) therapy.
    - Changing to a different TNF drug therapy, then one MRI (contrast as requested) of a single joint can be performed.
    - Addition of other treatments, including joint injections
- MRI or US should NOT be considered for routine follow-up of treatment.

## **Background and Supporting Information**

- \*Examples of appropriate laboratory studies may include: Lyme titers, rheumatoid factor (RF), anti-cyclic citrullinated peptide (anti-CCP), sedimentation rate (ESR), C-reactive protein (CRP), and antinuclear antibody (ANA)], joint fluid analysis

# Pigmented Villonodular Synovitis (PVNS) (MS-15.2)

---

MS.RA.0015.2.A

v1.0.2024

- MRI of the affected joint without contrast or CT of the affected joint with contrast (arthrogram) if MRI contraindicated is supported following plain x-rays.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider

## References (MS-15)

v1.0.2024

1. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Wrist Pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
2. Luchs JS, Flug JA, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Ankle Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
3. Hayes CW, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Elbow Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2022. <https://acsearch.acr.org/docs/69423/Narrative/>.
4. Jacobson JA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Extremity Joint Pain-Suspected Inflammatory Arthritis. *Am Coll Radiol (ACR)*; New: 2016. <https://acsearch.acr.org/docs/3097211/Narrative/>.
5. Wise JN, Weissman BN, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
6. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
7. Boutry N, Morel M, Flipo RM, et al. Early rheumatoid arthritis: a review of MRI and sonographic findings. *AJR Am J Roentgenol*. 2007;189:1502-1509. doi:10.2214/AJR.07.2548.
8. Murphey MD, Rhee JH, Lewis RB, et al. Pigmented villonodular synovitis: radiologic-pathologic correlation. *Radiographics*. 2008;28:1493-1518. doi:10.1148/rg.285085134.
9. Conaghan P, Edmonds J, Emery P, et al. Magnetic resonance imaging in rheumatoid arthritis: summary of OMERACT activities, current status, and plans. *Journal of Rheumatology*. 2001;28(5):1158-1161. <http://www.jrheum.org/content/28/5/1158.long>.
10. Ostergaard M, McQueen FM, Bird P, et al. Magnetic resonance imaging in rheumatoid arthritis--advances and research priorities. *Journal of Rheumatology*. 2005;32(12):2462-2464. <http://www.jrheum.org/content/32/12/2462.long>.
11. McQueen FM. The use of MRI in early RA. *Rheumatology*. 2008;47(11):1597-1599. doi:10.1093/rheumatology/ken332.
12. Gossec L, Fautrel B, Pham T, et al. Structural evaluation in the management of patients with rheumatoid arthritis: development of recommendations for clinical practice based on published evidence and expert opinion. *Joint Bone Spine*. 2005;72(3):229-234. doi:10.1016/j.jbspin.2004.10.011.
13. Cohen SB, Potter H, Deodhar A, et al. Extremity magnetic resonance imaging in rheumatoid arthritis: updated literature review. *Arthritis Care & Research*. 2011;63(5):660-665. doi:10.1002/acr.20413.
14. Singh JA, Furst DE, Bharat A, et al. 2012 update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care & Research*. 2012;64(5):625-639. doi:10.1002/acr.21641.
15. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis & Rheumatism (Arthritis Care & Research)*. 2008;59:762-784. doi:10.1002/art.23721.

# Post-Operative Joint Replacement Surgery (MS-16)

---

## Guideline

---

Post-Operative Joint Replacement Surgery – General (MS-16.1)  
References (MS-16)

# Post-Operative Joint Replacement Surgery – General (MS-16.1)

MS.PS.0016.1.A

v1.0.2024

- CT without contrast, MRI without contrast, or nuclear medicine studies (see: **MS-28** for nuclear medicine studies) with **ALL** of the following:
  - Recent plain x-ray is nondiagnostic
  - Suspected aseptic loosening of orthopaedic joint replacements
    - CT shoulder without contrast (CPT<sup>®</sup> 73200) can be performed as additional imaging following plain x-rays regardless of plain x-ray findings. See: **Shoulder (MS-19)**
      - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- CT without contrast or MRI without contrast with **ALL** of the following:
  - Negative plain x-ray
  - High suspicion for a periprosthetic fracture
    - CT Shoulder without contrast (CPT<sup>®</sup> 73200) can be performed as additional imaging following plain x-rays regardless of plain x-ray findings. See: **Shoulder (MS-19)**
      - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
- Joint aspiration is the initial evaluation after plain x-ray for a painful joint replacement when periprosthetic infection is suspected.
  - Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider
  - For suspected infection with negative or inconclusive joint aspiration culture see: **Nuclear Medicine (MS-28)**
- MRI Hip without contrast (CPT<sup>®</sup> 73721) or Ultrasound (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) are both appropriate for **EITHER** of the following:
  - Diagnosis of ALVAL (aseptic lymphocytic-dominated vasculitis-associated lesion) pseudotumors surrounding metal-on-metal (MoM) hip prostheses. One of these two imaging modalities can be approved but not both. See: **Soft Tissue Mass or Lesion of Bone (MS-10)**
  - Metal-On-Metal (MoM) Hip Prostheses that are considered high risk for implant performance issues from THA (Total hip arthroplasty) cup-neck impingement and subsequent ALTR (adverse local tissue reaction) with Co and Cr ion levels greater than 10 ppb.
- CT Hip without contrast (CPT<sup>®</sup> 73700) **OR** MRI Hip without contrast (CPT<sup>®</sup> 73721):

- Evaluate suspected particle disease (aggressive granulomatous disease) of the hip when infection has been excluded.
- For specific joints post-operative from replacement surgery:
  - See: **Shoulder (MS-19)**
  - See: **Elbow (MS-20)**
  - See: **Hip (MS-24)**
  - See: **Knee (MS-25)**
  - See: **Ankle (MS-26)**

### ***Background and Supporting Information***

- Complications following joint replacement surgery include (not limited to) periprosthetic fracture, infection, aseptic loosening, failure of fixation/component malposition, and wear.



## References (MS-16)

v1.0.2024

1. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
2. Hochman MG, Melenevsky YV, Metter DF, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Imaging After Total Knee Arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69430/Narrative/>.
3. Gyftopoulos S, Rosenberg ZS, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Imaging After Shoulder Arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2021. <https://acsearch.acr.org/docs/3097049/Narrative/>.
4. Weissman BN, Palestro CJ, Fox MG, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Imaging After Total Hip Arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2023. Available at: <https://acsearch.acr.org/docs/3094200/Narrative>
5. Toms AD, Davidson D, Masri BA, et al. Management of peri-prosthetic infection in total joint arthroplasty. *J Bone Joint Surg Br.* 2006;88(2):149-155. doi:10.1302/0301-620X.88B2.17058.
6. Love C, Marwin SE, Tomas MB, et al. Diagnosing infection in the failed joint replacement: A comparison of coincidence detection 18F-FDG and 111In-labeled leukocyte/99mTc-sulfur colloid marrow imaging. *J Nucl Med.* 2004;45(11):1864-1871.
7. Nawabi DH, Gold S, Lyman SL, et al. MRI predicts ALVAL and tissue damage in metal-on-metal hip arthroplasty. *Clin Orthop Relat Res.* 2014;472(2):471-481. doi:10.1007/s11999-013-2788-y.
8. Verberne SJ, Raijmakers PG, Temmerman OP. The accuracy of imaging techniques in the assessment of periprosthetic hip infection: a systematic review and meta-analysis. *J Bone Joint Surg Am.* 2016;98(19):1638-45. doi:10.2106/JBJS.15.00898.
9. Fritz J, Meshram P, Stern SE, Fritz B, Srikumaran U, McFarland EG. Diagnostic performance of advanced metal artifact reduction MRI for periprosthetic shoulder infection. *J Bone Joint Surg Am.* 2022;104:1352-1361. doi:10.2106/JBJS.21.00912.

# Limb Length Discrepancy (MS-17)

---

## Guideline

---

Limb Length Discrepancy (MS-17.1)  
Reference (MS-17)

## Limb Length Discrepancy (MS-17.1)

MS.LL.0017.1.A

v1.0.2024

- Either plain radiographic or “CT scanogram,” both reported with CPT<sup>®</sup> 77073, is appropriate to radiographically evaluate limb length discrepancy due to congenital anomalies, acquired deformities, growth plate (physeal injuries or surgery), or inborn errors of metabolism.
  - A diagnostic advanced imaging CPT code (e.g., CPT<sup>®</sup> 73700, CPT<sup>®</sup> 73701, or CPT<sup>®</sup> 73702) is not indicated for evaluation of limb length discrepancy.

## Reference (MS-17)

**v1.0.2024**

1. Leitzes A, Potter HG, Amaral T, et al. Reliability and accuracy of MRI scanogram in the evaluation of limb length discrepancy. *J Pediatr Orthop*. 2005;25(6):747-749.

# Anatomical Area Tables – General Information (MS-18)

---

## Guideline

---

Anatomical Area Tables – General Information (MS-18)

# Anatomical Area Tables – General Information (MS-18)

MS.AA.0018.A

v1.0.2024

The imaging guidelines for each anatomical area are presented in table format. The table below includes a description of how each column header should be utilized for each guideline **Shoulder (MS-19)** through **Foot (MS-27)**.

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in <a href="#">General Guidelines (MS-1.0)</a> )			
<b>Condition</b> (Individual's condition)	<b>Conservative Treatment</b> (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	<b>Advanced Imaging</b> (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	<b>Comments</b> (Additional comments related to the condition.)

# Shoulder (MS-19)

---

## Guideline

---

Shoulder (MS-19)  
References (MS-19)

# Shoulder (MS-19)

MS.SH.0019.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in [General Guidelines \(MS-1.0\)](#)

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Shoulder Pain</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>• US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) OR</li> <li>• CT Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73201) if MRI contraindicated</li> </ul>	
<b>Symptomatic Loose Bodies</b>	No	<ul style="list-style-type: none"> <li>• MRI Shoulder without contrast (CPT<sup>®</sup> 73221)</li> </ul>	
<b>Impingement</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>• MRI Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73222) OR</li> <li>• US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> <li>• CT Shoulder with contrast (CPT<sup>®</sup> 73201) if MRI is contraindicated</li> </ul>	



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)**

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Tendonitis/ Bursitis</b>	Yes		<ul style="list-style-type: none"> <li>• MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>• US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>
<b>Tendon Rupture (Biceps Long Head)</b>	No		<ul style="list-style-type: none"> <li>• When clinical exam is inconclusive due to inability to visualize a “Popeye” sign clinically, or for preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>
<b>Tendon Rupture (Pectoralis Major/Minor)</b>	No		<ul style="list-style-type: none"> <li>• When clinical exam is inconclusive, or for preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ MRI Chest without contrast (CPT<sup>®</sup> 71550) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Shoulder Rotator Cuff Tear (Complete and Partial)</b>	Yes*	<ul style="list-style-type: none"> <li>• MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>• MRI Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73222) OR</li> <li>• US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) OR</li> <li>• CT Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73201) if MRI is contraindicated</li> </ul>	<p>*Conservative treatment is not required with an acute shoulder injury prior to the onset of symptoms and consideration of surgery.</p> <p>If surgery is being considered, MRI without contrast, MRI with contrast (arthrogram), or CT arthrogram are required per <b><u>Shoulder Surgery- Arthroscopic and Open Procedure (CMM-315)</u></b></p>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)**

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Partial Tendon Rupture (Excluding Partial Rotator Cuff Tears)</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/ muscle tears.
<b>Complete Rupture – Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Shoulder Labral Tear (e.g., SLAP, ALPSA, HAGL)</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73222) OR</li> <li>• MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>• CT Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73201)</li> </ul>	For surgery criteria, see: <b><u>Shoulder Surgery- Arthroscopic and Open Procedures (CMM-315)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)**

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<p><b>Shoulder Dislocation/ Subluxation/ Instability, or Bankart/ Hill-Sachs Lesions</b></p>	<p>Yes*</p>	<ul style="list-style-type: none"> <li>• Individuals 40 years of age or younger with a first time dislocation, and in individuals with recurrent dislocations, conservative treatment not required:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73222) or</li> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221)</li> </ul> </li> <li>• CT Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73201) or CT Shoulder without contrast (CPT<sup>®</sup> 73200) if MRI is contraindicated</li> </ul>	<p>*Conservative treatment is required in individuals over age 40 with a first time dislocation.</p> <p>For surgery criteria, see: <b><u>Shoulder Surgery- Arthroscopic and Open Procedures (CMM-315)</u></b></p>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)**

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Frozen Shoulder/ Adhesive Capsulitis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Shoulder without contrast (CPT<sup>®</sup> 73221)</li> </ul>	For surgery criteria, see: <b><u>Manipulation Under Anesthesia (CMM-310)</u></b> and <b><u>Shoulder Surgery- Arthroscopic (CMM-315)</u></b>
<b>Avascular Necrosis (AVN) of the Humeral Head</b>	No	<ul style="list-style-type: none"> <li>See: <b><u>AVN (MS-4.1)</u></b></li> </ul>	
<b>Acromio-clavicular (AC) Separation</b>	No	<ul style="list-style-type: none"> <li>MRI Shoulder without contrast (CPT<sup>®</sup> 73221) to rule out possible rotator cuff tear following AC separation</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)**

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Sterno-clavicular (SC) Dislocation</b>	No	<ul style="list-style-type: none"> <li>• CT Chest without contrast (CPT<sup>®</sup> 71250) if posterior SC dislocation is evident or suspected</li> </ul>	
<b>Post-Operative Shoulder Surgery for Impingement, Rotator Cuff Tear, and/or Labral Tear</b>	Yes	<ul style="list-style-type: none"> <li>• In symptomatic individuals:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ MRI Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73222)</li> </ul> </li> <li>• US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) is also appropriate in symptomatic individuals following rotator cuff repair</li> <li>• CT Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73201) if MRI contraindicated</li> </ul>	

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<p><b>Preoperative Shoulder (Glenohumeral) Replacement Surgery</b></p>	<p>Yes</p>	<ul style="list-style-type: none"> <li>CT Shoulder without contrast (CPT<sup>®</sup> 73200) and/or MRI Shoulder without contrast (CPT<sup>®</sup> 73221) for preoperative planning prior to shoulder replacement</li> </ul>	<p>See also:  <u><b>Osteoarthritis (MS-12)</b></u>                      For joint surgery criteria, see:  <u><b>Shoulder Arthroplasty/ Arthrodesis (CMM-318)</b></u></p>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)**

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Post-Operative Shoulder (Glenohumeral) Replacement Surgery</b>	No	<ul style="list-style-type: none"> <li>• For suspected aseptic loosening or fracture as additional imaging following plain x-rays:                             <ul style="list-style-type: none"> <li>◦ CT Shoulder without contrast (CPT<sup>®</sup> 73200) OR</li> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) OR</li> <li>◦ Bone scan (CPT<sup>®</sup> 78315) OR</li> <li>◦ Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803 or CPT<sup>®</sup> 78831) OR</li> </ul> </li> </ul>	See also: <b><u>Post-Operative Joint Replacement (MS-16)</u></b>



After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<ul style="list-style-type: none"> <li>◦ Hybrid SPECT/CT (CPT<sup>®</sup> 78830)</li> <li>• For suspected infection with negative or inconclusive joint aspiration culture:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73321) OR</li> <li>◦ MRI Shoulder without and with contrast (CPT<sup>®</sup> 73223) OR</li> <li>◦ CT Shoulder with contrast (CPT<sup>®</sup> 73201) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) OR</li> <li>◦ See also: <b><u>Nuclear Medicine (MS-28)</u></b></li> </ul> </li> <li>• For possible rotator cuff tear:</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated as described in General Guidelines (MS-1.0)**

Condition (Individual's Condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<ul style="list-style-type: none"> <li>◦ CT Shoulder with contrast (arthrogram) (CPT<sup>®</sup> 73201) OR</li> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> <li>• For possible nerve injury:                             <ul style="list-style-type: none"> <li>◦ MRI Shoulder without contrast (CPT<sup>®</sup> 73221) OR</li> <li>◦ US Shoulder (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	

## References (MS-19)

v1.0.2024

1. Amini B, Beckmann NM, Beaman FD, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Traumatic. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69433/Narrative/>.
2. Neviasser RJ, Neviasser TJ. Recurrent instability of the shoulder after age 40. *J Shoulder Elbow Surg.* 1995;4(6):416-418.
3. Bradley M, Tung G, Green A. Overutilization of shoulder magnetic resonance imaging as a diagnostic screening tool in patients with chronic shoulder pain. *J Shoulder Elbow Surgery.* 2005;14(3):233-237. doi:10.1016/j.jse.2004.08.002.
4. Fongemie AE, Buss DD, and Rolnick SJ. Management of shoulder impingement syndrome and rotator cuff tears. *Am Fam Physician.* 1998;57(4):667-674. <https://www.aafp.org/afp/1998/0215/p667.html>.
5. Griffin LY. *Essentials of Musculoskeletal Care.* 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:212.
6. Gyftopoulos S, Rosenberg ZS, Roberts CC, ET. Al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Shoulder Arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/3097049/Narrative/>.
7. Hovelius L, Olofsson A, Sandstrom B, et al. Nonoperative treatment of primary anterior shoulder dislocation in patients forty years of age and younger: a prospective twenty-five year follow-up. *J Bone Joint Surg.* 2008;90:945-52. doi:10.2106/JBJS.G.00070.
8. Lin A, Gasbarro G, Sakr M. Clinical Applications of Ultrasonography in the Shoulder and Elbow. *J Am Acad Orthop Surg.* 2018;26:303-312.
9. Magee T. 3-T MRI of the shoulder: is MR arthrography necessary? *AJR J Am Roentgenol.* 2009;192:86-92. doi:10.2214/AJR.08.1097.
10. Major NM, Browne J, Domzalski T, Cothran RL, Helms CA. Evaluation of the glenoid labrum with 3-T MRI: is intraarticular contrast necessary. *AJR Am J Roentgenol.* 2011;196:1139-1144. doi:10.2214/AJR.08.1734.
11. McDonald LS, Dewing CB, Shupe PG, et al. Disorders of the proximal and distal aspects of the biceps muscle. *J Bone Joint Surg.* 2013;95:1235-1245. doi:10.2106/JBJS.L.00221.
12. Petersen SA, Murphy TP. The timing of rotator cuff repair for the restoration of function. *Journal of Shoulder and Elbow Surgery.* 2011;20(1):62-68. doi:10.1016/j.jse.2010.04.045.
13. Rehman A, Robinson P. Sonographic evaluation of injuries of the pectoralis muscles. *AJR Am J Roentgenol.* 2005;184:1205-1211. doi:10.2214/ajr.184.4.01841205.
14. Small KM, Adler RS, Shah SH, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Atraumatic. *Am Coll Radiol (ACR)*; New 2018. <https://acsearch.acr.org/docs/3101482/Narrative/>.
15. Steinbach LS, Chung CB, Yoshioka H. Technical Considerations for MRI of Upper Extremity Joints. In: Chung CB, Steinbach LS, eds. *MRI of the Upper Extremity Shoulder, Elbow, Wrist and Hand* Philadelphia, PA: Lippincott Williams & Wilkins 2010:211.
16. Streubel PN, Krych AJ, Simone JP, et al. Anterior glenohumeral instability: a pathology-based surgical treatment strategy. *J Am Acad Orthop Surg.* 2014;22:283-294. doi:10.5435/JAAOS-22-05-283.
17. Werner BC, Brockmeier SF, and Miller MD. Etiology, diagnosis, and management of failed SLAP repair. *J Am Acad Orthop Surg.* 2014;22(9):554-565. doi:10.5435/JAAOS-22-09-554.
18. Woodward TW and Best TM. The painful shoulder: Part II. Acute and chronic disorders. *Am Fam Physician.* 2000;61(11):3291-3300. <https://www.aafp.org/afp/2000/0601/p3291.html>.
19. Zappia M, Di Pietto F, Aliprandi A, et al. Multi-modal imaging of adhesive capsulitis of the shoulder. *Insights Imaging.* 2016;7:365-371.
20. Frankle MA, Teramoto A, Luo Z-P, Levy JC, Pupello D. Glenoid morphology in reverse shoulder arthroplasty: Classification and surgical implications. *Journal of Shoulder and Elbow Surgery.* 2009;18(6):874-885. doi:10.1016/j.jse.2009.02.013.

21. Beaman FD, von Herrmann PF, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected osteomyelitis, septic arthritis, or soft tissue infection (excluding spine and diabetic foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022.
22. Kowalczuk M, Elmaraghy A. Pectoralis major rupture: Evaluation and management. *J Am Acad Orthop Surg*. 2022;30:e617-e627. doi:10.5435/JAAOS-D-21-00541.
23. Fritz J, Meshram P, Stern SE, Fritz B, Srikumaran U, McFarland EG. Diagnostic performance of advanced metal artifact reduction MRI for periprosthetic shoulder infection. *J Bone Joint Surg Am*. 2022;104:1352-1361. doi:10.2106/JBJS.21.00912.

# Elbow (MS-20)

---

## Guideline

---

[Elbow \(MS-20\)](#)

[References \(MS-20\)](#)

# Elbow (MS-20)

MS.EB.0020.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Elbow Pain</b>	Yes	<ul style="list-style-type: none"> <li>MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Symptomatic Loose Bodies</b>	No	<ul style="list-style-type: none"> <li>MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>MRI Elbow with contrast (arthrogram) (CPT<sup>®</sup> 73222) <b>OR</b></li> <li>CT Elbow without contrast (CPT<sup>®</sup> 73200) <b>OR</b></li> <li>CT Elbow with contrast (arthrogram) (CPT<sup>®</sup> 73201)</li> </ul>	
<b>Tendonitis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Bursitis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Elbow without and with contrast (CPT<sup>®</sup> 73223) <b>OR</b></li> <li>MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Lateral (tennis elbow) or Medial (golfer's elbow) Epicondylitis</b>	Yes	<ul style="list-style-type: none"> <li>• To confirm clinical diagnosis of epicondylitis if symptoms persist for longer than 6 months despite at least 6 weeks conservative treatment in the last 3 months:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221)</li> <li><b>OR</b></li> <li>◦ US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	Epicondylitis, caused by tendon degeneration and tear of the common extensor tendon laterally or of the common flexor tendon medially, is a common clinical diagnosis for which imaging is not medically necessary except as noted.

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Suspected Osteochondral Injury</b>	No	<ul style="list-style-type: none"> <li>• If plain x-rays are negative and an osteochondral fracture is still suspected:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ MRI Elbow with contrast (arthrogram) (CPT<sup>®</sup> 73222) <b>OR</b></li> <li>◦ CT Elbow without contrast (CPT<sup>®</sup> 73200) <b>OR</b></li> <li>◦ CT Elbow with contrast (arthrogram) (CPT<sup>®</sup> 73201)</li> </ul> </li> </ul>	See: <b><u>Chondral/ Osteochondral Lesions (MS-13)</u></b> for other osteochondral injury scenarios
<b>Ruptured Biceps Insertion at Elbow</b>	No	<ul style="list-style-type: none"> <li>• When clinical exam is inconclusive or for preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Ruptured Triceps Insertion at Elbow</b>	No	<ul style="list-style-type: none"> <li>• When clinical exam is inconclusive or for preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/muscle tears.
<b>Complete Rupture – Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Trauma</b>	No	<ul style="list-style-type: none"> <li>• When surgery is being considered:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ CT Elbow without contrast (CPT<sup>®</sup> 73200)</li> </ul> </li> </ul>	
<b>Ulnar Collateral Ligament (UCL) Tear</b>	No	<ul style="list-style-type: none"> <li>• Following acute or repetitive (including overhead throwing athletes) elbow trauma:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow with contrast (arthrogram) (CPT<sup>®</sup> 73222) <b>OR</b></li> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) <b>OR</b></li> <li>◦ CT Elbow with contrast (arthrogram) (CPT<sup>®</sup> 73201)</li> </ul> </li> </ul>	
<b>Suspected Nerve Abnormality</b>	NA	<ul style="list-style-type: none"> <li>• This condition is imaged according to the criteria found in the Peripheral Nerve Disorder Guidelines. See: <b>Focal Neuropathy (PN-2)</b> in the Peripheral Nerve Disorders Imaging Guidelines</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>• CT Elbow without contrast (CPT<sup>®</sup> 73200) in symptomatic post-operative individuals following surgical treatment of complex fractures; <b>OR</b></li> <li>• MRI Elbow without contrast (CPT<sup>®</sup> 73221) in symptomatic post-operative individuals following soft-tissue surgery</li> </ul>	
<b>Preoperative Elbow Replacement Surgery</b>	Yes	<ul style="list-style-type: none"> <li>• CT Elbow without contrast (CPT<sup>®</sup> 73200) for preoperative planning prior to elbow replacement when congenital or post-traumatic deformities exist</li> </ul>	See: <b><u>Osteoarthritis (MS-12)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Post-Operative Elbow Replacement Surgery</b>	No	<ul style="list-style-type: none"> <li>• For suspected aseptic loosening or periprosthetic fracture when recent plain x-ray is nondiagnostic:                             <ul style="list-style-type: none"> <li>◦ CT Elbow without contrast (CPT<sup>®</sup> 73200) <b>OR</b></li> <li>◦ Bone scan (CPT<sup>®</sup> 78315) <b>OR</b></li> <li>◦ Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803 or 78831) <b>OR</b></li> <li>◦ Hybrid SPECT/CT (CPT<sup>®</sup> 78830)</li> </ul> </li> <li>• For suspected infection with negative or inconclusive joint aspiration culture:                             <ul style="list-style-type: none"> <li>◦ MRI Elbow without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ MRI Elbow without and with contrast (CPT<sup>®</sup> 73223) <b>OR</b></li> <li>◦ CT Elbow with contrast (CPT<sup>®</sup> 73201) <b>OR</b></li> <li>◦ US Elbow (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) <b>OR</b></li> <li>◦ See also: <b>Nuclear Medicine (MS-28)</b></li> </ul> </li> </ul>	

## References (MS-20)

**v1.0.2024**

1. McDonald LS, Dewing CB, Shupe PG, et al. Disorders of the proximal and distal aspects of the biceps muscle. *J Bone Joint Surg*. 2013;95:1235-1245. doi:10.2106/JBJS.L.00221.
2. Torp-Pedersen TE, Torp-Pedersen ST, Qvistgaard E, et al. Effect of glucocorticosteroid injections in tennis elbow verified on colour Doppler ultrasonography: evidence of inflammation. *Br J Sports Med*. 2008;42(12):978-982. doi:10.1136/bjsm.2007.041285.
3. Johnson GW, Cadwallader K, Scheffel SB, et al. Treatment of lateral epicondylitis. *Am Fam Physician*. 2007;76(6):843-848. <https://www.aafp.org/afp/2007/0915/p843.html>.
4. Griffin LY. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:279-280.
5. Hayes CW, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Elbow Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2022. <https://acsearch.acr.org/docs/69423/Narrative/>.
6. Bruce JR, Andrews JR. Ulnar collateral ligament injuries in the throwing athlete. *J Am Acad Orthop Surg*. 2014;22:315-325.
7. Beltran J, Rosenberg ZS. Diagnosis of compressive and entrapment neuropathies of the upper extremity: value of MR imaging. *AJR Am J Roentgenol*. 1994;163(3):525-531. doi:10.2214/ajr.163.3.8079837.
8. Lin A, Gasbarro G, Sakr M. Clinical Applications of Ultrasonography in the Shoulder and Elbow. *J Am Acad Orthop Surg*. 2018;26:303-312.
9. Stanborough RO, Wessell DE, Elhassan BT, Schoch BS. MRI of the elbow: Interpretation of common orthopaedic injuries. *J Am Acad Orthop Surg*. 2022;30:e573-e583. doi:10.5435/JAAOS-D-21-00193.
10. Beaman FD, von Herrmann PF, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/3094201/Narrative/>.

# Wrist (MS-21)

---

## Guideline

---

Wrist (MS-21)

References (MS-21)

# Wrist (MS-21)

MS.WR.0021.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Wrist Pain</b>	Yes		<ul style="list-style-type: none"> <li>• MRI Wrist without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>• US Wrist (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>
<b>Tendonitis</b>	Yes		<ul style="list-style-type: none"> <li>• MRI Wrist without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>• US Wrist (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Kienbock's Disease (Avascular Necrosis (AVN) of the Lunate)/ Preiser's Disease (Avascular Necrosis (AVN) of the Scaphoid)</b>	No	See <b>AVN (MS-4.1)</b>	
<b>Suspected Navicular/ Scaphoid Fracture</b>	No	<ul style="list-style-type: none"> <li>• When suspected based on history and physical exam:                             <ul style="list-style-type: none"> <li>◦ MRI Wrist without contrast (CPT<sup>®</sup> 73221)</li> <li><b>OR</b></li> <li>◦ CT Wrist without contrast (CPT<sup>®</sup> 73200)</li> </ul> </li> </ul>	See also: <b><u>Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints (MS-5.2)</u></b>



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Distal Radioulnar Joint (DRUJ) Instability</b>	No	<ul style="list-style-type: none"> <li>CT of both wrists without contrast (CPT<sup>®</sup> 73200) (should include wrists in supination and pronation)</li> </ul>	
<b>Complex Distal Radius/ Ulna Fracture</b>	No	<ul style="list-style-type: none"> <li>CT Wrist without contrast (CPT<sup>®</sup> 73200)</li> </ul>	
<b>Carpal Tunnel Syndrome/ Ulnar Tunnel Syndrome</b>	NA	<ul style="list-style-type: none"> <li>This condition is imaged according to the criteria found in the Peripheral Nerve Disorder Guidelines. See <b>Focal Neuropathy (PN-2)</b> in the Peripheral Nerve Disorders Imaging Guidelines</li> </ul>	
<b>Intrinsic Ligament (e.g. scapholunate)/ Triangular Fibrocartilage Complex (TFCC) Injuries</b>	Yes	<ul style="list-style-type: none"> <li>MRI Wrist with contrast (arthrogram) (CPT<sup>®</sup> 73222) <b>OR</b></li> <li>CT Wrist with contrast (arthrogram) (CPT<sup>®</sup> 73201)</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Complete Rupture - Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Wrist without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ US Wrist (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Wrist without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ US Wrist (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/muscle tears.
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>• CT Wrist without contrast (CPT<sup>®</sup> 73200) in symptomatic individuals following surgery for navicular/scaphoid fractures and complex distal radius/ulna fractures <b>OR</b></li> <li>• MRI Wrist with contrast (arthrogram) (CPT<sup>®</sup> 73222) in symptomatic individuals following DRUJ or TFCC surgery</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Preoperative Wrist Replacement Surgery</b>	Yes	<ul style="list-style-type: none"> <li>CT Wrist without contrast (CPT<sup>®</sup> 73200) for preoperative planning prior to wrist replacement when congenital or post-traumatic deformities exist</li> </ul>	See also: <b><u>Osteoarthritis (MS-12)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Post-Operative Wrist Replacement Surgery</b>	No	<ul style="list-style-type: none"> <li>• For suspected aseptic loosening or periprosthetic fracture when recent plain x-ray is nondiagnostic:                             <ul style="list-style-type: none"> <li>◦ CT Wrist without contrast (CPT<sup>®</sup> 73200) <b>OR</b></li> <li>◦ Bone scan (CPT<sup>®</sup> 78315, 78300, or 78306) <b>OR</b></li> <li>◦ Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803 or CPT<sup>®</sup> 78831) <b>OR</b></li> <li>◦ Hybrid SPECT/CT (CPT<sup>®</sup> 78830)</li> </ul> </li> <li>• For suspected infection with negative or inconclusive joint aspiration culture:                             <ul style="list-style-type: none"> <li>◦ MRI Wrist without contrast (CPT<sup>®</sup> 73221) <b>OR</b></li> <li>◦ MRI Wrist without and with contrast (CPT<sup>®</sup> 73223) <b>OR</b></li> <li>◦ CT Wrist with contrast (CPT<sup>®</sup> 73201) <b>OR</b></li> <li>◦ US Wrist (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) <b>OR</b></li> <li>◦ See also: <b>Nuclear Medicine (MS-28)</b></li> </ul> </li> </ul>	

**One Study/Area Only**

In hand and wrist advanced imaging, studies are frequently ordered of both areas. This is unnecessary since wrist MRI will image from above the wrist to the mid-metacarpal area. **Only one CPT<sup>®</sup> code should be reported.**

## References (MS-21)

v1.0.2024

1. Bruno MA, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hand and Wrist Trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
2. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Wrist Pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
3. Hayter CL, Gold SL, Potter HG. Magnetic resonance imaging of the wrist: bone and cartilage injury. *J Magn Reson Imaging*. 2013;37(5):1005-19. doi:10.1002/jmri.23845.
4. Pruitt DL, Gilula LA, Manske PR, et al. Computed tomography scanning with image reconstruction in evaluation of distal radius fractures. *J Hand Surg Am*. 1994;19(5):720-727. doi:10.1016/0363-5023(94)90174-0.
5. Magee T. Comparison of 3-T MRI and arthroscopy of intrinsic wrist ligament and TFCC tears. *AJR Am J Roentgenol*. 2009;192:80-85. doi:10.2214/AJR.08.1089.
6. Lee RK, Ng AW, Tong CS, et al. Intrinsic ligament and triangular fibrocartilage complex tears of the wrist: comparison of MDCT arthrography, conventional 3-T MRI, and MR arthrography. *Skeletal Radiol*. 2013;42:1277-85. doi:10.1007/s00256-013-1666-8.
7. Pahwa S, Srivastava DN, Sharma R, et al. Comparison of conventional MRI and MR arthrography in the evaluation wrist ligament tears: A preliminary experience. *Indian J Radiol Imaging*. 2014;3:259-67. doi:10.4103/0971-3026.137038.
8. Beaman FD, von Herrmann PF, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/3094201/Narrative/>.

# Hand (MS-22)

---

## Guideline

---

Hand (MS-22)

References (MS-22)

# Hand (MS-22)

MS.HA.0022.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Hand Pain</b>	Yes	<ul style="list-style-type: none"> <li>MRI Hand or Finger without contrast (CPT<sup>®</sup> 73218)</li> <li><b>OR</b></li> <li>US Hand (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Tendonitis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Hand or Finger without contrast (CPT<sup>®</sup> 73218)</li> <li><b>OR</b></li> <li>US Hand or Finger (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Occult Fracture</b>	No	<ul style="list-style-type: none"> <li>Advanced imaging guided by: <b><u>Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints (MS-5.2)</u></b></li> </ul>	
<b>Complex Fracture</b>	No	<ul style="list-style-type: none"> <li>CT Hand or Finger without contrast (CPT<sup>®</sup> 73200) when plain x-ray shows a complex fracture</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Ulnar Collateral Ligament (UCL) Thumb Injury</b>	No	<ul style="list-style-type: none"> <li>• If rule out for Stener lesion or complete tear of UCL of the thumb MCP joint:                             <ul style="list-style-type: none"> <li>◦ MRI Thumb without contrast (CPT<sup>®</sup> 73218)</li> <li><b>OR</b></li> <li>◦ US Thumb (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	Also called "Gamekeeper's Thumb" or "Skier's Thumb"
<b>Complete Rupture – Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Hand or Finger without contrast (CPT<sup>®</sup> 73218) <b>OR</b></li> <li>◦ US Hand or Finger (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Hand or Finger without contrast (CPT<sup>®</sup> 73218) <b>OR</b></li> <li>◦ US Hand or Finger (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/muscle tears.
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>• In symptomatic post-operative individuals following surgical treatment for complex hand or finger fractures or following soft-tissue surgery:                             <ul style="list-style-type: none"> <li>◦ CT Hand or Finger without contrast (CPT<sup>®</sup> 73200) <b>OR</b></li> <li>◦ MRI Hand or Finger without contrast (CPT<sup>®</sup> 73218)</li> </ul> </li> </ul>	

### **One Study/Area Only**

In hand and wrist advanced imaging, studies are frequently ordered of both areas. This is unnecessary since wrist MRI will image from above the wrist to the mid-metacarpal area. **Only one CPT<sup>®</sup> code should be reported.**

---

## References (MS-22)

---

**v1.0.2024**

1. Bruno MA, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hand and Wrist Trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
2. Hayter CL, Gold SL, Potter HG. Magnetic resonance imaging of the wrist: Bone and cartilage injury. *J Magn Reson Imaging*. 2013;37(5):1005-19. doi:10.1002/jmri.23845.

# Pelvis (MS-23)

---

## Guideline

---

[Pelvis \(MS-23\)](#)

[References \(MS-23\)](#)

# Pelvis (MS-23)

MS.PE.0023.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Pain-Pelvis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Pelvis without contrast (CPT<sup>®</sup> 72195) <b>OR</b></li> <li>MRI RT and/or LT Hip without contrast (CPT<sup>®</sup> 73721)</li> </ul>	
<b>Tendonitis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Pelvis without contrast (CPT<sup>®</sup> 72195) <b>OR</b></li> <li>MRI RT and/or LT Hip without contrast (CPT<sup>®</sup> 73721)</li> </ul>	
<b>Occult/ Insufficiency Fracture</b>	No	<ul style="list-style-type: none"> <li>MRI Pelvis without contrast (CPT<sup>®</sup> 72195) <b>OR</b></li> <li>CT Pelvis without contrast (CPT<sup>®</sup> 72192)</li> </ul>	See also: <b><u>Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints (MS-5.2)</u></b> for occult and stress fractures of the pelvis

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Complex Fracture/ Dislocation - Pelvis, Sacrum and Acetabulum</b>	No	<ul style="list-style-type: none"> <li>• CT Pelvis without contrast (CPT<sup>®</sup> 72192)</li> </ul>	Additionally, 3D rendering may be appropriate for preoperative planning.  See: <b>3D Rendering (MS-3)</b>
<b>Sacro-iliac (SI) Joint Pain, Sacroiliitis, Coccydynia</b>	Yes	<ul style="list-style-type: none"> <li>• Advanced imaging guided by:                             <ul style="list-style-type: none"> <li>◦ <b>Sacroiliac (SI) Joint Pain/ Sacroiliitis (SP-10.1)</b> in the Spine Imaging Guidelines</li> <li>◦ <b>Coccydynia without Neurological Features (SP-5.2)</b> in the Spine Imaging Guidelines</li> </ul> </li> </ul>	
<b>Piriformis Syndrome</b>	NA	<ul style="list-style-type: none"> <li>• This condition is imaged according to the criteria found in the Peripheral Nerve Disorder Guidelines. See: <b>Focal Neuropathy (PN-2)</b> in the Peripheral Nerve Disorders Imaging Guidelines</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>MRI Pelvis without contrast (CPT<sup>®</sup> 72195) for a suspected partial tendon rupture of a specific named tendon not otherwise specified</li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/muscle tears.
<b>Osteitis Pubis/ Symphysis Pubis Diastasis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Pelvis without contrast (CPT<sup>®</sup> 72195)</li> </ul>	
<b>Athletic Pubalgia (Sports Hernia)</b>	Yes	<ul style="list-style-type: none"> <li>To evaluate for the cause of suspected athletic pubalgia:                             <ul style="list-style-type: none"> <li>MRI Pelvis without contrast (athletic pubalgia protocol) (CPT<sup>®</sup> 72195) <b>OR</b></li> <li>Dynamic pelvic ultrasound (CPT<sup>®</sup> 76857)</li> </ul> </li> </ul>	
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>CT Pelvis without contrast (CPT<sup>®</sup> 72192) in symptomatic individuals following surgery for complex pelvic ring/acetabular fractures</li> </ul>	

## References (MS-23)

**v1.0.2024**

1. Bencardino JT, Stone TJ, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/> .
2. Mehta S, Auerbach JD, Born CT, et al. Sacral fractures. *J Am Acad Orthop Surg*. 2006;14:656-665.
3. Omar IM, Zoga AC, Kavanagh EC, et al. Athletic Pubalgia and "Sports Hernia": Optimal MR Imaging Technique and Findings. *RadioGraphics*. 2008;28:1415-1438. doi:10.1148/rg.285075217.
4. Khan W, Zoga AC, Meyers WC. Magnetic Resonance Imaging of Athletic Pubalgia and the Sports Hernia - Current Understanding and Practice. *Magn Reson Imaging Clin N Am*. 2013;21:97-110. doi:10.1016/j.mric.2012.09.008.
5. Morley N, Grant T, Blount K, et al. Sonographic evaluation of athletic pubalgia. *Skeletal Radiol*. 2016 May;45(5):689-99. doi:10.1007/s00256-016-2340-8.
6. Caudill P, Nyland J, Smith C, et al. Sports hernias: a systematic literature review. *British Journal of Sports Medicine*. 2008;42(12):954-964. doi:10.1136/bjism.2008.047373.
7. Suarez JC, Ely EE, Mutnal AB, et al. Comprehensive approach to the evaluation of groin pain. *Journal of the American Academy of Orthopaedic Surgeons*. 2013;21:558-570. doi:10.5435/JAAOS-21-09-558.
8. Heer ST, Callander JW, Kraeutler MJ, Mei-Dan O, Mulcahey MK. Hamstring Injuries. *The Journal of Bone and Joint Surgery*. 2019;101(9):843-853. doi:10.2106/jbjs.18.00261.



# Hip (MS-24)

---

## Guideline

---

Hip (MS-24)

References (MS-24)

# Hip (MS-24)

MS.HI.0024.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Hip Pain</b>	Yes		<ul style="list-style-type: none"> <li>• MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• US Hip (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>
<b>Symptomatic Loose Bodies</b>	No		<ul style="list-style-type: none"> <li>• MRI Hip without contrast (CPT<sup>®</sup> 73721)</li> </ul>
<b>Tendonitis/ Bursitis</b>	Yes		<ul style="list-style-type: none"> <li>• MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• US Hip (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>
<b>Hip Abductor Tendon Tear/ Avulsion</b>	No		<ul style="list-style-type: none"> <li>• MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• US Hip (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Complete Rupture – Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Hip (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Hip (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/ muscle tears.

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Occult/ Insufficiency Fracture</b>	No	<ul style="list-style-type: none"> <li>• MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• CT Hip without contrast (CPT<sup>®</sup> 73700)</li> </ul>	See also: <b><u>Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints (MS-5.2)</u></b> for occult and stress fractures of the hip
<b>Avascular Necrosis (AVN) of the Femoral Head</b>	No	<ul style="list-style-type: none"> <li>• See: <b><u>AVN (MS-4.1)</u></b></li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
Labral Tear	Yes	<ul style="list-style-type: none"> <li>• MRI Hip with contrast (arthrogram) (CPT<sup>®</sup> 73722)</li> <li><b>OR</b></li> <li>• CT Hip with contrast (arthrogram) (CPT<sup>®</sup> 73701)</li> <li><b>OR</b></li> <li>• MRI Hip without contrast (CPT<sup>®</sup> 73721)</li> </ul>	For surgery criteria, see: <b><u>Hip Surgery-Arthroscopic and Open Procedures (CMM-314)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Femoroacetabular Impingement</b>	Yes	<ul style="list-style-type: none"> <li>• For preoperative planning for femoroacetabular impingement:                             <ul style="list-style-type: none"> <li>◦ MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ MRI Hip with contrast (arthrogram) (CPT<sup>®</sup> 73722)</li> </ul> </li> </ul> <p>IN ADDITION TO:</p> <ul style="list-style-type: none"> <li>• CT Hip without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>• CT Pelvis without contrast (CPT<sup>®</sup> 72192)</li> </ul>	For surgery criteria, see: <b><u>Hip Surgery-Arthroscopic and Open Procedures (CMM-314)</u></b>
<b>Piriformis Syndrome</b>	NA	<ul style="list-style-type: none"> <li>• This condition is imaged according to the criteria found in the Peripheral Nerve Disorder Guidelines. See <b>Focal Neuropathy (PN-2)</b> in the Peripheral Nerve Disorders Imaging Guidelines</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>• Symptomatic individuals following surgery for labral tears and femoroacetabular impingement:                             <ul style="list-style-type: none"> <li>◦ MRI Hip with contrast (arthrogram) (CPT<sup>®</sup> 73722)</li> </ul> </li> <li>• Symptomatic individuals following surgery for hip fracture and/or hip avascular necrosis:                             <ul style="list-style-type: none"> <li>◦ CT Hip without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>◦ MRI Hip without contrast (CPT<sup>®</sup> 73721)</li> </ul> </li> </ul>	
<b>Preoperative Hip Replacement Surgery</b>	Yes	<ul style="list-style-type: none"> <li>• CT Hip without contrast (CPT<sup>®</sup> 73700) or CT Pelvis without contrast (CPT<sup>®</sup> 72192) for preoperative planning prior to hip replacement when congenital or post-traumatic deformities exist</li> </ul>	See also: <b><u>Osteoarthritis (MS-12)</u></b>  For surgery criteria, see: <b><u>Hip Arthroplasty- Total and Partial (CMM-313)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<p><b>Post-Operative Hip Replacement Surgery</b></p>	<p>No*</p>	<ul style="list-style-type: none"> <li>• For suspected aseptic loosening of hip replacement when recent plain x-ray is nondiagnostic:                             <ul style="list-style-type: none"> <li>◦ CT Hip without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>◦ Bone scan (CPT<sup>®</sup> 78315) <b>OR</b></li> <li>◦ Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803 or CPT<sup>®</sup> 78831) <b>OR</b></li> <li>◦ Hybrid SPECT/CT (CPT<sup>®</sup> 78830)</li> </ul> </li> <li>• For suspected infection with negative or inconclusive joint aspiration culture:                             <ul style="list-style-type: none"> <li>◦ MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> </ul> </li> </ul>	<p>See: <b><u>Post-Operative Joint Replacement Surgery (MS-16)</u></b></p>



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<ul style="list-style-type: none"> <li>◦ MRI Hip without and with contrast (CPT<sup>®</sup> 73723) <b>OR</b></li> <li>◦ CT Hip with contrast (CPT<sup>®</sup> 73701) <b>OR</b></li> <li>◦ US Hip (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) <b>OR</b></li> <li>◦ See also: <b><u>Nuclear Medicine (MS-28)</u></b></li> <li>• For a suspicion of a periprosthetic fracture when recent plain x-ray is nondiagnostic:                         <ul style="list-style-type: none"> <li>◦ CT Hip without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>◦ Bone scan (CPT<sup>®</sup> 78315) <b>OR</b></li> <li>◦ Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup></li> </ul> </li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<p>78803 or CPT<sup>®</sup> 78831)  <b>OR</b></p> <ul style="list-style-type: none"> <li>◦ Hybrid SPECT/CT (CPT<sup>®</sup> 78830)</li> <li>• To evaluate component malposition or heterotopic bone after plain x-ray:                             <ul style="list-style-type: none"> <li>◦ CT Hip without contrast (CPT<sup>®</sup> 73700)</li> </ul> </li> <li>• For possible nerve injury:                             <ul style="list-style-type: none"> <li>◦ MRI Hip without contrast (CPT<sup>®</sup> 73721)</li> </ul> </li> <li>• For suspected tendinitis/ bursitis, abductor injury, or other soft tissue abnormality (*requires conservative treatment):                             <ul style="list-style-type: none"> <li>◦ MRI Hip without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Hip (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	

## References (MS-24)

v1.0.2024

1. Greene WB (Ed.). *Essentials of Musculoskeletal Care*. 2<sup>nd</sup> Ed. Rosemont, IL, American Academy of Orthopaedic Surgeons, 2001.
2. Manek NJ and Lane NE. Osteoarthritis: Current concepts in diagnosis and management. *Am Fam Physician*. 2000;61(6):1795-1804. <https://www.aafp.org/afp/2000/0315/p1795.html> .
3. Papadopoulos EC and Kahn SN. Piriformis syndrome and low back pain: a new classification and review of the literature. *Orthop Clin North Am*. 2004;35(1):65-71. doi:10.1016/S0030-5898(03)00105-6.
4. Reurink G, Sebastian, Bisselink JM, et al. Reliability and Validity of Diagnostic Acetabular Labral Lesions with Magnetic Resonance Arthrography. *J Bone Joint Surg A*. 2012;94(181):1643-1648. doi:10.2106/JBJS.K.01342.
5. Steinbach LS, Palmer WE, and Schweitzer ME. Special Focus Session MR Arthrography1. *RadioGraphics*. 2002;22(5):1223-1246.
6. Redmond JM, Chen AW, and Domb BG. Greater Trochanteric Pain Syndrome. *J Am Acad Orthop Surg*. 2016;24(4):231-240. doi:10.5435/JAAOS-D-14-00406.
7. Center for Devices and Radiological Health. Metal-on-Metal Hip Implants - Information for Orthopaedic Surgeons. U S Food and Drug Administration Home Page.
8. Ward RJ, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Acute Hip Pain-Suspected Fracture. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Revised: 2018. <https://acsearch.acr.org/docs/3082587/Narrative/>.
9. Weissman BN, Palestro CJ, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Imaging After Total Hip Arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2023. <https://acsearch.acr.org/docs/3094200/Narrative/>.
10. Mintz DN, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
11. Murphey MD, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Osteonecrosis of the Hip. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2022. <https://acsearch.acr.org/docs/69420/Narrative/> .
12. Bencardino JT, Stone TJ, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
13. Verberne SJ, Raijmakers PG, and Temmerman OP. The Accuracy of Imaging Techniques in the Assessment of Periprosthetic Hip Infection. A Systematic Review and Meta-Analysis. *J Bone Joint Surg Am*. 2016;98(19):1638-45. doi:10.2106/JBJS.15.00898.
14. Shin AY, Morin WD, Gorman JD, et al. The superiority of magnetic resonance imaging in differentiating the cause of hip pain in endurance athletes. *Am J Sports Med*. 1996;24:168-76. doi:10.1177/036354659602400209.
15. Slocum KA, Gorman JD, Puckett ML, et al. Resolution of abnormal MR signal intensity in patients with stress fractures of the femoral neck. *AJR Am J Roentgenol*. 1997;168:1295-9. doi:10.2214/ajr.168.5.9129429.
16. Lee EY, Margherita AJ, Gierada DS, et al. MRI of Piriformis Syndrome. *American Journal of Roentgenology*. 2004;183:63-64. doi:10.2214/ajr.183.1.1830063.
17. Jankiewicz JJ, Hennrikus WL, and Houkom JA. The appearance of the piriformis muscle syndrome in computed tomography and magnetic resonance imaging: a case report and review of the literature. *Clin Orthop*. 1991;262:205-09.
18. Rossi P, Cardinali P, Serrao M, et al.. Magnetic resonance imaging findings in piriformis syndrome: a case report. *Arch Phys Med Rehabil*. 2001;82(4):519-21. doi:10.1053/apmr.2001.21971.
19. Heer ST, Callander JW, Kraeutler MJ, Mei-Dan O, Mulcahey MK. Hamstring Injuries. *The Journal of Bone and Joint Surgery*. 2019;101(9):843-853. doi:10.2106/jbjs.18.00261.
20. Beaman FD, von Herrmann PF, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine

and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/3094201/Narrative/>.

# Knee (MS-25)

---

## Guideline

---

Knee (MS-25)

References (MS-25)

# Knee (MS-25)

MS.KN.0025.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Knee Pain</b>	Yes	<ul style="list-style-type: none"> <li>MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>US Knee (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Symptomatic Loose Bodies</b>	No	<ul style="list-style-type: none"> <li>MRI Knee without contrast (CPT<sup>®</sup> 73721)</li> <li>CT Knee with contrast (arthrogram) (CPT<sup>®</sup> 73701) if MRI cannot be performed</li> </ul>	
<b>Tendonitis</b>	Yes	<ul style="list-style-type: none"> <li>MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>US Knee (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Complex Knee Fracture</b>	No	<ul style="list-style-type: none"> <li>MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>CT Knee without contrast (CPT<sup>®</sup> 73700)</li> </ul>	See also: <b><u>Fractures (MS-5)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Meniscus Tear</b>	Yes*	<ul style="list-style-type: none"> <li>• MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• CT Knee with contrast (arthrogram) (CPT<sup>®</sup> 73701) if MRI cannot be performed</li> </ul> <p>*Conservative treatment is not required if at least 2 of following 4 criteria are met:</p> <ol style="list-style-type: none"> <li>1) Positive McMurray's, positive Thessaly, or positive Apley's Compression Test</li> <li>2) twisting or acute injury of the knee</li> <li>3) locked knee/inability to fully extend the knee on exam in comparison to the opposite knee</li> <li>4) knee effusion</li> </ol>	For surgery criteria, see: <b><u>Knee Surgery-Arthroscopic and Open Procedures (CMM-312)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<ul style="list-style-type: none"> <li>MRI Knee without contrast (CPT<sup>®</sup> 73721) for clinical suspicion of a symptomatic degenerative meniscus tear in an individual with osteoarthritis following conservative treatment</li> </ul>	



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Ligament Tear</b>	Yes*	<ul style="list-style-type: none"> <li>• MRI Knee without contrast (CPT<sup>®</sup> 73721)</li> </ul> <p>*Conservative treatment is not required if any of the following signs are positive in comparison to the opposite knee:</p> <ul style="list-style-type: none"> <li>• Anterior drawer</li> <li>• Lachman</li> <li>• Pivot shift</li> <li>• Posterior drawer</li> <li>• Posterior sag</li> <li>• Valgus stress</li> <li>• Varus stress</li> </ul>	For surgery criteria, see: <b><u>Knee Surgery-Arthroscopic and Open Procedures (CMM-312)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Knee Joint Dislocation</b>	No	<ul style="list-style-type: none"> <li>• Following significant trauma to evaluate for ligament and vascular injury:                             <ul style="list-style-type: none"> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721) AND <i>EITHER</i></li> <li>◦ MR Angiography lower extremity without and with contrast (CPT<sup>®</sup> 73725) <b>OR</b></li> <li>◦ CT Angiography lower extremity without and with contrast (CPT<sup>®</sup> 73706)</li> </ul> </li> </ul>	
<b>Patellar Dislocation/ Subluxation</b>	No	<ul style="list-style-type: none"> <li>• MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b> CT Knee without contrast (CPT<sup>®</sup> 73700) when there is an acute knee injury, consideration of surgery, <b>AND</b> concern for osteochondral fracture or loose osteochondral fracture fragment</li> </ul>	For surgery criteria, see: <b><u>Knee Surgery- Arthroscopic and Open Procedures (CMM-312)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Recurrent Patellar Instability</b>	Yes	<ul style="list-style-type: none"> <li>MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b> CT Knee without contrast (CPT<sup>®</sup> 73700) if consideration for surgery</li> </ul>	For surgery criteria, see: <b><u>Knee Surgery-Arthroscopic and Open Procedures (CMM-312)</u></b>
<b>Patellofemoral Pain Syndrome/ Anterior Knee Pain/ Tracking Disorder</b>	Yes	<ul style="list-style-type: none"> <li>MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b> CT Knee without contrast (CPT<sup>®</sup> 73700) if consideration for surgery</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Suspected Osteochondral Injury</b>	No	<ul style="list-style-type: none"> <li>• If plain x-rays are negative and an osteochondral fracture is still suspected:                             <ul style="list-style-type: none"> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ MRI Knee with contrast (arthrogram) (CPT<sup>®</sup> 73722) <b>OR</b></li> <li>◦ CT Knee with contrast (arthrogram) (CPT<sup>®</sup> 73701)</li> </ul> </li> </ul>	See: <b><u>Chondral/ Osteochondral Lesions (MS-13)</u></b> for other osteochondral injury scenarios.  For surgery criteria, see: <b><u>Knee Surgery- Arthroscopic and Open Procedures (CMM-312)</u></b>
<b>Avascular Necrosis (AVN) of the Distal Femur</b>	No	<ul style="list-style-type: none"> <li>• See: <b><u>AVN (MS-4.1)</u></b></li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Baker's Cyst (Popliteal Cyst)</b>	Yes	<ul style="list-style-type: none"> <li>• US Knee (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) is the initial imaging study</li> <li>• MRI Knee without contrast (CPT<sup>®</sup> 73721) for preoperative planning</li> </ul>	See also: <b>Acute Limb Swelling (PVD-12)</b> in the Peripheral Vascular Disease Imaging Guidelines
<b>Plica (Symptomatic Synovial Plica/ Medial Synovial Shelf)</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Knee without contrast (CPT<sup>®</sup> 73721)</li> </ul>	
<b>Hemarthrosis</b>	No	<ul style="list-style-type: none"> <li>• MRI Knee without contrast (CPT<sup>®</sup> 73721) for clinical suspicion of cruciate ligament tear (requires a positive objective sign for ACL/PCL tear) or patellar dislocation (requires a positive apprehension sign)</li> <li>• CT Knee without contrast (CPT<sup>®</sup> 73700) for clinical suspicion of non-displaced intra-articular fracture</li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Complete Rupture of the Distal Quadriceps Tendon or Patellar Ligament/ Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Knee (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Knee (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/ muscle tears.

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Complete Rupture – Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Knee (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>• In symptomatic individuals following surgery for meniscus tears and reconstruction of the anterior cruciate ligament:                             <ul style="list-style-type: none"> <li>◦ MRI Knee with contrast (arthrogram) (CPT<sup>®</sup> 73722) <b>OR</b></li> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721)</li> </ul> </li> <li>• In symptomatic individuals following surgery for fracture/dislocation:                             <ul style="list-style-type: none"> <li>◦ CT Knee without contrast (CPT<sup>®</sup> 73700)</li> </ul> </li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Preoperative Knee Replacement Surgery</b>	Yes	<ul style="list-style-type: none"> <li>CT Knee without contrast (CPT<sup>®</sup> 73700) for preoperative planning prior to knee replacement when congenital or post-traumatic deformities exist of the patella, distal femur and/or proximal tibia</li> </ul>	See also: <u><b>Osteoarthritis (MS-12)</b></u>  For surgery criteria, see: <u><b>Knee Arthroplasty- Total and Partial (CMM-311)</b></u>



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
Post-Operative Knee Replacement Surgery	No*	<ul style="list-style-type: none"> <li>• For suspected aseptic loosening when recent plain x-ray is nondiagnostic:                             <ul style="list-style-type: none"> <li>◦ CT Knee without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ See also: <b><u>Nuclear Medicine (MS-28)</u></b></li> </ul> </li> <li>• For suspected infection with negative or inconclusive joint aspiration culture:                             <ul style="list-style-type: none"> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ MRI Knee without and with contrast (CPT<sup>®</sup> 73723) <b>OR</b></li> <li>◦ CT Knee with contrast (CPT<sup>®</sup> 73701) <b>OR</b></li> </ul> </li> </ul>	See also: <b><u>Post-Operative Joint Replacement Surgery (MS-16)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<ul style="list-style-type: none"> <li>◦ US Knee (CPT<sup>®</sup> 76881 or 76882)</li> <li>◦ See also: <b>Nuclear Medicine (MS-28)</b></li> <li>• Following plain x-ray for suspected periprosthetic fracture:                             <ul style="list-style-type: none"> <li>◦ CT Knee without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721)</li> <li>◦ 3-phase bone scan (CPT<sup>®</sup> 78315)</li> </ul> </li> <li>• For suspected osteolysis or component instability, rotation, or wear:                             <ul style="list-style-type: none"> <li>◦ CT Knee without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721)</li> </ul> </li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<ul style="list-style-type: none"> <li>• For suspected periprosthetic soft tissue abnormality unrelated to infection (e.g., tendinopathy, arthrofibrosis, patellar clunk syndrome, impingement of nerves or other soft tissue) *requires conservative treatment:                             <ul style="list-style-type: none"> <li>◦ MRI Knee without contrast (CPT<sup>®</sup> 73721)</li> <li><b>OR</b></li> <li>◦ US Knee (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	

## References (MS-25)

v1.0.2024

1. Harrison BK, Abell BE, Gibson TW. The Thessaly test for detection of meniscal tears: validation of a new physical examination technique for primary care medicine. *Clin J Sport Med*. 2009;19:9-12. doi:10.1097/JSM.0b013e31818f1689.
2. Landewé RBM, Günther KP, Lukas C, et al. EULAR/EFFORT recommendations for the diagnosis and initial management of patients with acute or recent onset swelling of the knee. *Ann Rheum Dis*. 2010;69:12-19. doi:10.1136/ard.2008.104406.
3. Johnson MW. Acute knee effusions: a systematic approach to diagnosis. *Am Fam Physician*. 2000;61(8):2391-2400. <https://www.aafp.org/afp/2000/0415/p2391.html>.
4. ACR Appropriateness Criteria, Nontraumatic knee pain, 2008.
5. Sung-Jae Kim, Byoung-Yoon Hwang, Choi DH, et al. *J Bone Joint Surg A*. 2012;94(16):e118 1-7.
6. Kannus P, Järvinen M. Nonoperative treatment of acute knee ligament injuries. A review with special reference to indications and methods. *Sports Med*. 1990;9(4):244-260. doi:10.2165/00007256-199009040-00005.
7. Manek NJ and Lane NE. Osteoarthritis: Current concepts in diagnosis and management. *Am Fam Physician*. 2000;61(6):1795-1804. <https://www.aafp.org/afp/2000/0315/p1795.html>.
8. Griffin LY. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:84, 541-545.
9. Lee IS, Choi JA, Kim TK, et al. Reliability analysis of 16-MDCT in preoperative evaluation of total knee arthroplasty and comparison with intraoperative measurements. *Am J Roentgenol*. 2006;186(6):1778-1782. doi:10.2214/AJR.05.1191.
10. Morrissey RT, Weinstein SL (Eds.). *Lovell and Winter's Pediatric Orthopaedics*. 6th Ed. Philadelphia, PA: Lippincott Williams and Wilkins; 2005:1413.
11. Woolson ST, Harris AHS, Wagner DW, et al; Component alignment during total knee arthroplasty with use of standard or custom instrumentation: A Randomized Clinical Trial Using Computed Tomography for Postoperative Alignment Measurement. *Journal of Bone and Joint Surgery*. 2014;96:366-372. doi:10.2106/JBJS.L.01722.
12. Vance K, Meredick R, Schweitzer ME, et al. Magnetic resonance imaging of the postoperative meniscus. *Arthroscopy*. 2009;25:522-30. doi:10.1016/j.arthro.2008.08.013.
13. Magee T, Shapiro M, and Williams D. Prevalence of meniscal radial tears of the knee revealed by MRI after surgery. *Am J Roentgenol*. 2004;184:931-936. doi:10.2214/ajr.182.4.1820931.
14. Meyers AB, Haims AH, Menn K, et al. Imaging of anterior cruciate ligament repair and its complications. *Am J Roentgenol*. 2010;194:476-484. doi:10.2214/AJR.09.3200.
15. Tuite, MJ, Kransdorf MJ, Beaman FD, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Acute Trauma to the Knee. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2019. <https://acsearch.acr.org/docs/69419/Narrative>.
16. Bennett DL, Nelson JW, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Nontraumatic Knee Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
17. Zoga AC, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Soft-Tissue Masses. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
18. Hochman MG, Melenevsky YV, Metter DF, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria<sup>®</sup> Imaging After Total Knee Arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2023. Available at: <https://acsearch.acr.org/docs/69430/Narrative/>.
19. Kurosaka M, Yagi M, Yoshiya S, Muratsu H, Mizuno K. Efficacy of the axially loaded pivot shift test for the diagnosis of a meniscal tear. *Int Orthop*. 1999;23:271-274. doi:10.1007/s002640050369.
20. Fowler PJ, Lubliner JA. The predictive value of five clinical signs in the evaluation of meniscal pathology. *Arthroscopy*. 1989;5(3):184-186. doi:10.1016/0749-8063(89)90168-0.

21. Beaman FD, von Herrmann PF, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/3094201/Narrative/>.
22. Pauyo T, Park JP, Bozzo I, Bernstein M. Patellofemoral instability part I: evaluation and nonsurgical treatment. *J Am Acad Orthop Surg.* 2022;30:e1431-e1442. doi:10.5435/JAAOS-D-22-00254.

# Ankle (MS-26)

---

## Guideline

---

Ankle (MS-26)

References (MS-26)

# Ankle (MS-26)

MS.AL.0026.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Ankle Pain</b>	Yes		<ul style="list-style-type: none"> <li>• MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• US Ankle (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>
<b>Symptomatic Loose Bodies</b>	No		<ul style="list-style-type: none"> <li>• MRI Ankle without contrast (CPT<sup>®</sup> 73721)</li> </ul>
<b>Complex Fracture</b>	No		<ul style="list-style-type: none"> <li>• MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• CT Ankle without contrast (CPT<sup>®</sup> 73700)</li> </ul>
<b>Ankle Sprain, Including Avulsion Fracture</b>	Yes		<ul style="list-style-type: none"> <li>• MRI Ankle Without Contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• CT Ankle without contrast (CPT<sup>®</sup> 73700)</li> </ul>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>High Ankle Sprain (Syndesmosis Injury)</b>	No	<ul style="list-style-type: none"> <li>• MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>• CT Ankle without contrast (CPT<sup>®</sup> 73700)</li> </ul>	
<b>Suspected Osteochondral Injury</b>	No	<ul style="list-style-type: none"> <li>• If plain x-rays are negative and an osteochondral fracture is still suspected, ONE of the following:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ CT Ankle without contrast (CPT<sup>®</sup> 73700)</li> </ul> </li> </ul>	See: <b><u>Chondral/Osteochondral Lesions (MS-13)</u></b> for other osteochondral injury scenarios
<b>Avascular Necrosis (AVN) of the Talus</b>	No	<ul style="list-style-type: none"> <li>• See: <b><u>AVN (MS-4.1)</u></b></li> </ul>	



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Anterior Impingement</b> <b>Anterior-Lateral Impingement</b> <b>Posterior Impingement (e.g., Os Trigonum Syndrome)</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Ankle with contrast (arthrogram) (CPT<sup>®</sup> 73722) <b>OR</b></li> <li>• CT Ankle with contrast (arthrogram) (CPT<sup>®</sup> 73701) <b>OR</b></li> <li>• MRI Ankle without contrast (CPT<sup>®</sup> 73721)</li> </ul>	
<b>Tendonitis</b>	Yes	<ul style="list-style-type: none"> <li>• For suspected posterior tibial dysfunction, peroneal tendon or subluxation, Achilles tendonitis:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Ankle (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Complete Rupture of Achilles Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative evaluation:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Ankle (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Complete Rupture -Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Ankle (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Ankle (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/ muscle tears.
<b>Instability</b>	Yes	<ul style="list-style-type: none"> <li>• For preoperative evaluation:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ MRI Ankle with contrast (arthrogram) (CPT<sup>®</sup> 73722)</li> </ul> </li> </ul>	
<b>Charcot Ankle</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Ankle without contrast (CPT<sup>®</sup> 73721)</li> </ul>	

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>• In symptomatic individuals following surgery for ligament/tendon injuries, <i>one of the following</i>:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ US Ankle (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> <li>• For symptomatic individuals following surgery for complex fractures:                             <ul style="list-style-type: none"> <li>◦ CT Ankle without contrast (CPT<sup>®</sup> 73700)</li> </ul> </li> </ul>	
<b>Preoperative Ankle Replacement Surgery</b>	Yes	<ul style="list-style-type: none"> <li>• CT Ankle without contrast (CPT<sup>®</sup> 73700) for preoperative planning prior to ankle replacement when congenital or post-traumatic deformities exist</li> </ul>	See: <b><u>Osteoarthritis (MS-12)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Post-Operative Ankle Replacement Surgery</b>	No	<ul style="list-style-type: none"> <li>• For suspected aseptic loosening or periprosthetic fracture when recent plain x-ray is nondiagnostic:                             <ul style="list-style-type: none"> <li>◦ CT Ankle without contrast (CPT<sup>®</sup> 73700) <b>OR</b></li> <li>◦ Bone scan (CPT<sup>®</sup> 78315, 78300, or 78306) <b>OR</b></li> <li>◦ Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803 or 78831) <b>OR</b></li> <li>◦ Hybrid SPECT/CT (CPT<sup>®</sup> 78830)</li> </ul> </li> <li>• For suspected infection with negative or inconclusive joint aspiration culture:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> </ul> </li> </ul>	See: <b><u>Post-Operative Joint Replacement Surgery (MS-16)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced Imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
		<ul style="list-style-type: none"> <li>◦ MRI Ankle without and with contrast (CPT<sup>®</sup> 73723) <b>OR</b></li> <li>◦ CT Ankle with contrast (CPT<sup>®</sup> 73701) <b>OR</b></li> <li>◦ US Ankle (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882) <b>OR</b></li> <li>◦ See also: <b><u>Nuclear Medicine (MS-28)</u></b></li> </ul>	

**One Study/Area Only**

In foot and ankle advanced imaging, studies are frequently ordered of both areas. This is unnecessary since ankle MRI will image from above the ankle to the mid-metatarsal area. **Only one CPT<sup>®</sup> code should be reported.**

## References (MS-26)

v1.0.2024

1. Donovan A, Rosenberg ZS. MRI of ankle and lateral hindfoot impingement syndromes. *AJR*. 2010;195:595-604.
2. Wolfe MW, Uhl TL, and McClusky LC. Management of ankle sprains. *Am Fam Physician* 2001;63(1):93-104. <https://www.aafp.org/afp/2001/0101/p93.html>.
3. Griffin LY. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:593-596; 606-609; 683.
4. Bergkvist D, Astrom I, Josefsson PO, et al. Acute Achilles Tendon Rupture: A Questionnaire Follow-up of 487 Patients. *J Bone Joint Surg Am*. 2012;94(13):1229-1233. doi:10.2106/JBJS.J.01601.
5. Hartgerink P, Fessell DP, Jacobson JA, et al. Full- versus partial-thickness Achilles tendon tears: sonographic accuracy and characterization in 26 cases with surgical correlation. *Radiology* 2001;220:406-412. doi: 10.1148/radiology.220.2.r01au41406.
6. Jones MP, Riaz JK, Smith RLC. Surgical Interventions for Treating Acute Achilles Tendon Rupture: Key Findings from a Recent Cochrane Review. *J Bone Joint Surg Am*. 2012;94(12):e88 1-6. doi:10.2106/jbjs.j.01829.
7. Vaseenon T, Amendola A. Update on anterior ankle impingement. *Current Reviews in Musculoskeletal Medicine*. 2012;5:140-150. doi:10.1007/s12178-012-9117-z.
8. Talusan PG, Toy J, Perez J, Milewski MD, et al. Anterior ankle impingement: diagnosis and treatment. *J Am Acad Orthop Surg*. 2014;22:333-339. doi:10.5435/JAAOS-22-05-333.
9. Nault ML, Kocher MS, Micheli LJ. Os Trigonum Syndrome. *J Am Acad Orthop Surg*. 2014;22:545-553. doi:10.5435/JAAOS-22-09-545.
10. Peace KAL, Jillier JC, Hulme A, et al. MRI features of posterior ankle impingement syndrome in ballet dancers: a review 25 cases. *Clinical Radiology*. 2004;59:1025-1033. doi:10.1016/j.crad.2004.02.010.
11. J Kane and R Zell. Achilles Tendon Rupture. Physician Resource Center. *American Orthopaedic Foot & Ankle Society*. Last reviewed July 2015.
12. Garras DN, et al. MRI is Unnecessary for Diagnosing Acute Achilles Tendon Ruptures. *Clinical Orthopaedics and Related Research*. 2012;470:2268–2273 Retrospective Analysis with finding. doi:10.1007/s11999-012-2355-y.
13. Mosher TJ, Kransdorf MJ, Adler R, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Ankle. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Revised: 2020. <https://acsearch.acr.org/docs/69436/Narrative/>.
14. Luchs JS, Flug JA, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Ankle Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
15. Dodd A, Daniels TR. Charcot Neuroarthropathy of the Foot and Ankle. *J Bone Joint Surg Am*. 2018;100:696-711. doi:10.2106/JBJS.17.00785.
16. Beaman FD, von Herrmann PF, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/3094201/Narrative/>.

# Foot (MS-27)

---

## Guideline

---

Foot (MS-27)

References (MS-27)

# Foot (MS-27)

MS.FT.0027.A

v1.0.2024

After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>General Foot Pain</b>	Yes	• MRI Foot without contrast (CPT <sup>®</sup> 73718)	
<b>Complex Fractures</b>	No	• CT Foot without contrast (CPT <sup>®</sup> 73700)	
<b>Plantar Plate Disorders, Including Turf Toe Injuries</b>	Yes	• MRI Foot without contrast (CPT <sup>®</sup> 73718)	
<b>Sesamoid Disorders</b>	Yes	• MRI Foot without contrast (CPT <sup>®</sup> 73718) <b>OR</b> • CT Foot without contrast (CPT <sup>®</sup> 73700)	



**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Lisfranc Tarsometatarsal Fracture or Dislocation</b>	No	<ul style="list-style-type: none"> <li>• MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>• CT Foot without contrast (CPT<sup>®</sup> 73700)</li> </ul>	
<b>Tarsal Navicular Stress/Occult Fracture</b>	No	<ul style="list-style-type: none"> <li>• MRI Foot without contrast (CPT<sup>®</sup> 73718)</li> <li>• Tc-99m bone scan foot (CPT<sup>®</sup> 78315) if MRI cannot be performed</li> <li>• CT Foot without contrast (CPT<sup>®</sup> 73700) for follow-up of healing fractures</li> </ul>	See also: <b><u>Suspected Occult/ Stress/ In-sufficiency Fracture/ Stress Reaction and Shin Splints (MS-5.2)</u></b>

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in General Guidelines [MS-1.0])**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Avascular Necrosis (AVN) of the Tarsal Navicular (Kohler Disease) or Metatarsal Head (Frieberg's Infraction)</b>	No	<ul style="list-style-type: none"> <li>• See: <b>AVN (MS-4.1)</b></li> </ul>	
<b>Tendonitis</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>• US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Complete Rupture – Tear of a Specific Named Tendon</b>	No	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>◦ US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Partial Tendon Rupture</b>	No	<ul style="list-style-type: none"> <li>• For a suspected partial tendon rupture of a specific named tendon not otherwise specified:                             <ul style="list-style-type: none"> <li>◦ MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>◦ US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	MRI is <i>NOT</i> needed for muscle belly strains/muscle tears.
<b>Morton's Neuroma</b>	Yes	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>◦ MRI Foot without and with contrast (CPT<sup>®</sup> 73720) <b>OR</b></li> <li>◦ US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	
<b>Plantar Fasciitis</b>	Yes*	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>◦ US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	*Provider-directed conservative treatment must be for 6 months or more.

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Suspected Plantar Fascia Rupture or Tear</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>• US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul>	
<b>Diabetic Foot Infection</b>	No	<ul style="list-style-type: none"> <li>• For suspected osteomyelitis or soft tissue infection as a complement to plain x-ray (both plain x-ray and MRI are indicated):                             <ul style="list-style-type: none"> <li>◦ MRI Foot without and with contrast (CPT<sup>®</sup> 73720) <b>OR</b></li> <li>◦ MRI Foot without contrast (CPT<sup>®</sup> 73718)</li> </ul> </li> </ul>	See also: <b><u>Infection-General (MS-9.1)</u></b>
<b>Tarsal Tunnel Syndrome including Baxter's Neuropathy</b>	Yes	<ul style="list-style-type: none"> <li>• For preoperative planning if mass/lesion is suspected as etiology of entrapment:                             <ul style="list-style-type: none"> <li>◦ MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>◦ US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> </ul>	

**After an initial plain x-ray has been obtained, and results are available to the provider, the following advanced imaging is indicated (as described in [General Guidelines \[MS-1.0\]](#))**

Condition (Individual's condition)	Conservative Treatment (Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required?) (Yes or No)	Advanced imaging (The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.)	Comments (Additional comments related to the condition.)
<b>Tarsal Coalition</b>	Yes	<ul style="list-style-type: none"> <li>• For preoperative planning:                             <ul style="list-style-type: none"> <li>◦ MRI Ankle without contrast (CPT<sup>®</sup> 73721) <b>OR</b></li> <li>◦ CT Ankle without contrast (CPT<sup>®</sup> 73700)</li> </ul> </li> </ul>	
<b>Sinus Tarsi Syndrome</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Ankle without contrast (CPT<sup>®</sup> 73721) if diagnosis is unclear or for preoperative evaluation</li> </ul>	
<b>Charcot Foot</b>	Yes	<ul style="list-style-type: none"> <li>• MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>• MRI Foot without and with contrast (CPT<sup>®</sup> 73720)</li> </ul>	
<b>CRPS Type I</b>	Yes	<ul style="list-style-type: none"> <li>• Triple phase bone scan (CPT<sup>®</sup> 78315) <b>OR</b></li> <li>• MRI Foot without contrast (CPT<sup>®</sup> 73718)</li> </ul>	
<b>Post-Operative</b>	Yes	<ul style="list-style-type: none"> <li>• In symptomatic individuals following surgery for conditions including the tendons, ligaments, and plantar plate, ONE of the following:                             <ul style="list-style-type: none"> <li>◦ MRI Foot without contrast (CPT<sup>®</sup> 73718) <b>OR</b></li> <li>◦ US Foot (CPT<sup>®</sup> 76881 or CPT<sup>®</sup> 76882)</li> </ul> </li> <li>• In symptomatic individuals following surgery for complex fractures, sesamoid fractures, and subtalar arthrodesis:                             <ul style="list-style-type: none"> <li>◦ CT Foot without contrast (CPT<sup>®</sup> 73700)</li> </ul> </li> </ul>	

### **One Study/Area Only**

In foot and ankle advanced imaging, studies are frequently ordered of both areas. This is unnecessary since ankle MRI will image from above the ankle to the mid- metatarsal area. **Only one CPT<sup>®</sup> code should be reported.**

## References (MS-27)

v1.0.2024

1. Griffin LY. *Essentials of Musculoskeletal Care*. 3<sup>rd</sup> edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:619-622;667-671;681-684;697-699;700-702.
2. Needell S, Cutler J. Morton neuroma imaging. *eMedicine*, April 11, 2011
3. Morton's Neuroma. MDGuidelines™.
4. Berquist TH. *Radiology of the Foot and Ankle*. 2nd Ed. Philadelphia, Lippincott, 2000, pp.155-156.
5. Bouche R. Sinus Tarsi Syndrome. What is Sinus Tarsi Syndrome, Testing and Treatment. [http://www.aapm.org/sinus\\_tarsi\\_syndrome.html](http://www.aapm.org/sinus_tarsi_syndrome.html).
6. D Resnick. *Internal Derangements of Joints 2006: Imaging-Arthroscopic Correlation*. Washington, DC. Oct.31-Nov. 4, 2006.
7. Doty JF, Coughlin MJ. Metatarsophalangeal joint instability of the lesser toes and plantar plate deficiency. *J Am Acad Orthop Surg*. 2014;22(4):235-245. doi:10.5435/JAAOS-22-04-235.
8. Lareau CR, Sawyer GA, Wang JH, et al. Plantar and medial heel pain: diagnosis and management. *J Am Acad Orthop Surg*. 2014;22:372-380. doi:10.5435/JAAOS-22-06-372.
9. Sung, W, Weil L Jr, Weill LS Sr, et al. Diagnosis of plantar plate injury by magnetic resonance imaging with reference to Intraoperative findings. *Journal of Foot Ankle Surgery*. 2012;51(5):570-574. doi:10.1053/j.jfas.2012.05.009.
10. Bancroft LW, Kransdorf MJ, Adler R, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Foot. *Am Coll Radiol (ACR)*; Date of Origin: 2010. Revised: 2019. <https://acsearch.acr.org/docs/70546/Narrative/>.
11. Wise JN, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
12. Bencardino JT, Stone TJ, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
13. Kransdorf MJ, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2019. <https://acsearch.acr.org/docs/69340/Narrative/>.
14. Thomas JL, Christensen JC, Kravitz SR, et al. The Diagnosis and Treatment of Heel Pain: A Clinical Practice Guideline - Revision 2010. *J Foot Ankle Surg*. 2010;49:S1-S19. doi: 10.1053/j.jfas.2010.01.001.
15. Goff JD, Crawford R. Diagnosis and Treatment of Plantar Fasciitis. *Am Fam Physician*. 2011 Sep;84(6):676-682. <https://www.aafp.org/afp/2011/0915/p676.html>.
16. Baxter D, Pfeffer G. Treatment of chronic heel pain by surgical release of the first branch of the lateral plantar nerve. *Clin Orthop*. 1992;279:229-236.
17. Schepsis A, Leach R, Gorzyca J. Plantar fasciitis: etiology, treatment, surgical results, and review of the literature. *Clin Orthop* 1991;266:185-196.
18. Neufeld SK, Cerato R. Plantar fasciitis: evaluation and treatment. *J Am Acad Orthop Surg*. 2008;16:338-46. doi:10.5435/00124635-200806000-00006.
19. Dodd A, Daniels TR. Charcot Neuroarthropathy of the Foot and Ankle. *J Bone Joint Surg Am*. 2018;100:696-711. doi:10.2106/JBJS.17.00785.

# Nuclear Medicine (MS-28)

---

## Guideline

---

Nuclear Medicine (MS-28)

References (MS-28)



## Nuclear Medicine (MS-28)

MS.NM.0028.A

v1.0.2024

Results of plain x-rays performed after the current episode of symptoms started or changed need to be available to the requesting provider, unless otherwise specified below.

- SPECT scan may be approved for any of the indications for which a bone scan can be approved.
  - If the request is for CPT<sup>®</sup> 78300 and CPT<sup>®</sup> 78803, then only CPT<sup>®</sup> 78803 is to be approved if medical necessity is established.
  - If the request is for CPT<sup>®</sup> 78305 or CPT<sup>®</sup> 78306 and CPT<sup>®</sup> 78803, then two CPT<sup>®</sup> codes may be approved if medical necessity is established.
- Nuclear Medicine may be used in the evaluation of some musculoskeletal disorders, and other rare indications exist as well.
  - Evaluation of suspected aseptic loosening of orthopedic prostheses when recent plain x-ray is nondiagnostic:
    - Bone scan (CPT<sup>®</sup> 78315) **OR**
    - Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803, or 78831) **OR**
    - Hybrid SPECT/CT (CPT<sup>®</sup> 78830)
    - See also: **Post-Operative Joint Replacement Surgery (MS-16)** and anatomic tables
  - For detection of ischemic or infarcted regions in sickle cell disease:
    - Nuclear medicine bone marrow imaging (CPT<sup>®</sup> 78102, 78103,78104) **OR**
    - SPECT (CPT<sup>®</sup> 78803) **OR**
    - Hybrid SPECT/CT (CPT<sup>®</sup> 78830)
    - See also: **Modality General Considerations (PEDMS-1.3)**
  - Evaluation of complex regional pain syndrome or reflex sympathetic dystrophy, after failure of six weeks provider-directed conservative treatment (per **General Guidelines (MS-1.0)**):
    - Triple phase bone scan (CPT<sup>®</sup> 78315)
    - See: **Foot (MS-27)** for imaging criteria of CRPS of the foot
    - For interventional pain criteria see: **Regional Sympathetic Blocks (CMM-209)** and **Spinal Cord Stimulators (CMM-211)**
  - Evaluation of Paget's disease
    - Bone scan (CPT<sup>®</sup> codes: 78300, 78305, 78306) **OR**
    - Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803) **OR**
    - Hybrid SPECT/CT (CPT<sup>®</sup> 78830)

- See also: **Soft Tissue Mass or Lesion of Bone (MS-10)**
- Suspected fractures
  - If criteria per **Suspected Occult/Stress/Insufficiency Fracture/Stress Reaction and Shin Splints (MS-5.2)** are met, but MRI cannot be performed:
    - Tc-99m bone scan whole body (CPT<sup>®</sup> 78306) with SPECT of the area of interest (CPT<sup>®</sup> 78803) **OR**
    - Hybrid SPECT/CT (CPT<sup>®</sup> 78830) **OR**
    - Bone scan (CPT<sup>®</sup> 78315, 78305, or 78300)
  - Evaluation of suspected bone infection if MRI or CT cannot be done and when infection is multifocal, or when the infection is associated with orthopedic hardware or chronic bone alterations from trauma or surgery
    - FDG PET/CT (CPT<sup>®</sup> 78815 for multifocal infection, or CPT<sup>®</sup> 78811 for unifocal/limited area of interest) if MRI or CT is equivocal or cannot be done
      - At this time, FDG is the only indicated radiotracer for use with PET/CT in the imaging of musculoskeletal conditions.
    - Bone scan (CPT<sup>®</sup> 78315, 78300, 78305, or 78306) **OR**
    - Distribution Of Radiopharmaceutical Agent SPECT (CPT<sup>®</sup> 78803 or 78831) **OR**
    - Hybrid SPECT/CT (CPT<sup>®</sup> 78830 or 78832)
    - A labeled leukocyte scan (radiopharmaceutical inflammatory imaging - one of CPT<sup>®</sup> codes: 78800, 78801, 78802, or 78803) in concert with Tc-99m sulfur colloid marrow imaging (one of CPT<sup>®</sup> codes: 78102, 78103, or 78104)
    - See also: **Post-Operative Joint Replacement Surgery (MS-16)**
    - For specific joints post-operative from replacement surgery:
      - See: **Shoulder (MS-19)**
      - See: **Elbow (MS-20)**
      - See: **Hip (MS-24)**
      - See: **Knee (MS-25)**
      - See: **Ankle (MS-26)**

## References (MS-28)

v1.0.2024

1. Beaman FD, von Herrmann PF, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. Revised: 2022. <https://acsearch.acr.org/docs/3094201/Narrative/>.
2. Kransdorf MJ, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2019. <https://acsearch.acr.org/docs/69340/Narrative/>.
3. Wise JN, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Revised: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
4. Bencardino JT, Stone TJ, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
5. Hochman MG, Melenevsky YV, Metter DF, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Total Knee Arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2023. <https://acsearch.acr.org/docs/69430/Narrative/>.
6. Weissman BN, Palestro CJ, Fox MG, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Total Hip Arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2023. Available at: <https://acsearch.acr.org/docs/3094200/Narrative/>.

# Policy History and Instructions for Use

---

## Guideline

---

Policy History and Instructions for Use

# Policy History and Instructions for Use

## Policy History and Instructions for Use v2.0.2024

### Instructions for Use

This Medical Policy provides assistance in interpreting United HealthCare Services, Inc. standard benefit plans. When deciding coverage, the federal, state (Ohio Administrative Code [OAC]) or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state (OAC) or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state (OAC) or contractual requirements for benefit plan coverage govern.

Before using this policy, please check the federal, state (OAC) or contractual requirements for benefit plan coverage. United HealthCare Services, Inc. reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

United HealthCare Services, Inc. uses InterQual<sup>®</sup> for the primary medical/surgical criteria, and the American Society of Addiction Medicine (ASAM) for substance use, in administering health benefits. If InterQual<sup>®</sup> does not have applicable criteria, United HealthCare Services, Inc. may also use United HealthCare Services, Inc.'s Medical Policies, Coverage Determination Guidelines, and/or Utilization Review Guidelines that have been approved by the Ohio Department for Medicaid Services. The United HealthCare Services, Inc.'s Medical Policies, Coverage Determination Guidelines, and Utilization Review Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

### Policy History/Revision Information

Date	Summary of Changes
02/01/2024	Annual evidence-based updates
07/01/2024	Interim evidence-based updates